

Original Research Article

EFFECTIVENESS OF STANDARDIZED PATIENTS AND ROLE-PLAY SIMULATION METHODS IN IMPROVING SELF-CONFIDENCE AND THERAPEUTIC COMMUNICATION SKILLS OF NURSING STUDENTS

Bagus Dwi Cahyono ^{1*}, Ida Zuhroidah ¹, Mokh. Sujarwadi ¹

¹Jember University, Pasuruan Campus, Indonesia

*Correspondence:

Bagus Dwi Cahyono

Jember University, Pasuruan Campus, Indonesia

KH. Mansyur road 207, Tembokrejo Village, Purworejo Sub-District, Pasuruan City, East Java Province, Indonesia - 67118

Email: bagusdwi.akper@unej.ac.id

Article Info:

Received: 12 December 2019

Revised: 16 January 2020

27 January 2020

Accepted: 27 January 2020

DOI:

<https://doi.org/10.36720/nhjk.v9i1.128>

Abstract

Background: Communication skills are critical skills that must be possessed by nurses because communication is a dynamic process that is used to collect assessment data, provide education or health information to influence clients to apply it in life, show caring, provide comfort, foster self-confidence and, value respect - client value. Nurses in communicating with clients must have high confidence in the effect of these communications.

Objectives: The purpose of this study was to analyze the differences in the effect of standardized patient simulation methods and role play in increasing self-confidence and therapeutic communication skills of students of the Nursing Academy of Pasuruan.

Methods: The design of this study is the quasi experimental study. There are two therapeutic communication training groups. Group A uses a Standardized Patient (SP) and, Group B uses the role play (RP) method. The population in this study were students of the Nursing Academy of Pasuruan, and the sample was taken by simple random sampling with a sample size group of 16 standardized patients and a roleplay group of 16 people. The measuring instrument used was an observation sheet on therapeutic communication skills and a questionnaire on confidence in therapeutic communication. The effectiveness of the two methods was analyzed by the non-parametric test, namely the Mann Whitney test, to test 2 unpaired samples.

Results: The results of this study indicate that there are differences in standardized patient methods and role play on self-confidence and therapeutic communication skills in nursing academy students who are statistically significant at 0.03 ($p=0.05$).

Conclusion: The standardized patient method is more effective in increasing self-confidence and therapeutic communication skills in nursing academy students compared to the roleplay method.

Keywords: Standardized Patient, Roleplay, Therapeutic Communication, Self-Confidence

INTRODUCTION

Communication skills are an important professional element. The aim of the problem-based learning (PBL) learning model is to provide nurse students with the communication skills needed to assist the nursing process. Communication is a powerful therapeutic tool, and a nurse needs it to achieve health goals to influence others (Ustun, 2005). According to Ozkan, 1992; Tattersall and Ellis 1998 and Terakye, 1991 in Ustun, 2005 stated that patients and their families would feel uncomfortable communicating with nurses who are not in accordance with their level of knowledge, many patients find it difficult to explain the situation to nurses and do not understand what is explained by the nurse.

Communication skills are critical skills that must be possessed by nurses because communication is a dynamic process that is used to collect assessment data, provide education or health information to influence clients to apply it in life, show caring, provide comfort, foster self-confidence, and value respect - client value. Nurses who have therapeutic communication skills will not only easily establish a relationship of trust with clients, prevent legal problems, provide professional satisfaction in nursing services, and improve the image of the nursing profession and the image of the hospital (Suryani, 2014).

Effective communication is an important skill reported to be less encouraging to oncology nurses at the Tokyo Metropolitan Cancer Screening Center because it eliminates many opportunities to confirm difficult information, for example, when patient and family support is given poor information such as cancer diagnosis (Fukui *et al*, 2008). Results Survey reports of inpatient and family comfort in the Mardi Rahayu Hospital ER from 2006 to May 2009, there were still between 5% and 6.5% of respondents feeling uncomfortable when in the ER all because of inadequate communication according to the patient so that the respondent gave advice as well as criticism in providing services to provide explanations to

patients with friendliness and more to provide information with good and polite communication (Hermawan, 2010).

Simulations in nursing education in the form of static manikin and role play (Sanford, 2010). Standardized patient is a simulation which is a terminology used to simulate patients (trained as simulated patients who are sick) as actual patients. This is an expensive high-tech instrument. Standardized Patients are also used in formative and summative evaluations of communication skills (Bosse *et al*, 2010). As an experiential learning technique, simulations encourage students to connect concepts, apply theory to practice, and make clinical decisions (Waldner and Olson, 2007). The actual context of the simulation is to improve communication, teamwork, delegation, priority settings, and leadership skills Bauman *et al*, 2008). Reading and Webster (2013) found that simulations on the complexity of drug administration as a catalyst to improve error reports, which, in turn, improvised practices were affected by changes in procedures and policies. Aebersold *et al* (2012) incorporating simulations in the nursing education curriculum to explore critical thinking skills. Students are not only capable of their skills, but case studies provide a realistic context for developing decision-making skills.

Roleplay as a learning strategy that is suitable for the social family model (Joyce and Weil, 2000). This strategy emphasizes social learning naturally and sees cooperative behavior as a stimulus for students between social and intellectuals. This role-play strategy students represent and express the characters they know in everyday life (Blatner, 2009). The use of role-playing emphasizes personal attention, problems, behavior and active participation (Hober, 2012). In a literature review by Lanea and Rollnick (2007) explains that simulated patients and role-play are often used in learning communication skills in the world.

Blum *et al* (2010) explains that the conceptualization of confidence in a nurse as a symbol of trust and respect in the relationship between nurses and patients. Simulation has

been proven as a learning strategy that can increase clinical confidence and competence.

The purpose of this study was to analyze the differences in the effectiveness of standardized patients and role-play simulation methods in improving self-confidence and therapeutic communication skills of students at the Nursing Academy of Pasuruan.

METHODS

Study Design

The research method in this study was used Quasy-Experimental Study.

Setting

This research was conducted in the Nursing Academy of Pasuruan.

Research Subject

Samples in this study were selected using simple random sampling based on the formulation of Supranto (2000), namely:

$$(t-1)(r-1) \geq 15$$

Explanation:

t = Number of treatment groups

r = Number of replications

According that formulation, the sample size in this study was 16 respondents in each group (standardized patient and role play groups). The respondents obtained through the determination of the sample by drawing.

RESULTS

Characteristics of Respondents by Gender

Table 1. Distribution of Frequency of Respondents by Gender in the Nursing Academy of Pasuruan on 2018.

Gender	Role Play		Standardized Patient	
	Frequency (f)	Percentage (%)	Frequency (f)	Percentage (%)
Male	7	43.75	5	31.25
Female	9	56.25	11	68.75
Total	16	100.00	16	100.00

Sources: Primary Data of Questionnaire, 2018.

Based on the result above (table 1), it obtained that the gender of respondents was mostly female in each group, namely 9 respondents in the role play group (56.25%) and 11 respondents in the standardized patient group (68.75%).

Instruments

Data were collected using therapeutic communication skill observation sheet and the questionnaire of self-confidence. Therapeutic communication skill observation sheet consists of three domains, namely introducing (2 items), non-verbal communication (7 items), and verbal communication (2 items). The questionnaire of self-confidence consists of eight domains, namely optimistic (2 items), responsibility (1 item), positive thinking (3 items), independent (5 items), dare to rejection (1 item), calm (3 items), realistic (1 item), and relax (6 items). The reliability test (Cronbach Alpha) of the therapeutic communication skill observation sheet was 0.60. The reliability test (Cronbach Alpha) of the questionnaire of self-confidence was 0.849.

Data Analysis

The analysis method in this study was used non-parametric tests using Mann Whitney Test with two unpaired samples test.

Ethical Consideration

This research implemented based on permission from director of Nursing Academy of Pasuruan with permit letter number 800/Akper/398/423.104.10/2018.

Analysis of Effectiveness of the Self-Confidence and Therapeutic Communication Skills in the Role Play Method Group and Standardized Patient Method Group

Table 2. Analysis of Effectiveness of the Self-Confidence and Therapeutic Communication Skills in the Role Play Method Group and Standardized Patient Method Group using Mann-Whitney U Test in the Nursing Academy of Pasuruan on 2018.

Description	Self-Confidence		Therapeutic Communication Skills	
	Role Play	Standardized Patient	Role Play	Standardized Patient
Mean	68.43	76.38	64.86	68.69
SD	7.70	6.82	5.60	4.17
Min	48	65	52	59
Max	82	86	72	72
	<i>p</i> -value = 0.01		<i>p</i> -value = 0.03	

Sources: Primary Data of Questionnaire, 2018.

The results on the table 2 found that self-confidence in the standardized patient group (mean = 76.38; SD = 6.82; min = 65; max = 86) is better than role play group (mean = 68.43; SD = 7.70; min = 48; max = 82). In addition, the above results also showed that therapeutic communication skills in standardized patient group (mean = 68.69; SD = 4.17; min = 59; max = 72) is better than the role play group (mean = 64.86; SD = 5.60; min = 52; max = 72).

For the results of the analysis using the Mann-Whitney U Test obtained the difference between the two groups either in self-confidence (*p* = 0.01) and therapeutic communication skills (*p* = 0.03).

Based on the above results, it can be concluded that learning using the standardized patient method is more effective than the role play method for increasing self-confidence and therapeutic communication skills in nursing students.

DISCUSSION

Based on the results of the statistical analysis shows that the average value of the standardized patient method is higher than the role-play method and the statistical test results, there are significant differences (*p* = 0.05) between the standardized patient method and role play in increasing self-confidence and therapeutic communication skills. This means that the standardized patient method is more effective in increasing self-confidence and therapeutic communication skills to students compared to the role-play method. Standardized Patient is a type of simulation that trains patients or simulated actors to provide a history of the disease reliably and consistently (Bosse et al., 2010). Research conducted by Hans M Bosse, Martin Nickel et al. (2010) published by the journal BioMed Central Medical Education with the title "Peer role-play

and standardized patients in communication training: a comparative study on the student's perspective on acceptability, realism, and perceived effects." In the perceived effect of training in both groups (5.17 ± 0.37 for the role playgroup, 5.50 ± 0.43 for the standardized patient's group; 6 = strongly agree, 1 = strongly disagree) there were significant differences in the standardized patient's group (*p* < 0.003). While there is no significant difference between acceptability and realism between the two groups. The results of the study by Wenker (2009), explained that standardized patients provide positive experiences for physiotherapy students in the first year in conducting interviews with patients as measured by the Arizona Clinical Interview Rating Scale.

Standardized patient and role play are actually included in the simulation learning method which means as if dealing with the real

situation, the stages are orientation, training, implementation, and debriefing. The difference is if the standardized patient simulation uses patients or other people who are considered capable of playing the real patient. In this research, the standardized patient is a psychiatric nursing lecturer, wherein the curriculum of psychiatric nursing; there are competency demands that are capable of conducting therapeutic communication, whereas role play is done by playing the role of fellow friends.

Self-confidence is a positive attitude of an individual that enables him to develop positive assessments both towards oneself and the environment (Jasnita, 2002). Simulation has been proven as a learning strategy that can increase clinical confidence and competence (Blum *et al*, 2010).

Therapeutic communication is a way to foster a therapeutic relationship where information is conveyed and the exchange of feelings and thoughts to influence others, this communication is planned consciously aims, and activities are centered on healing the patient. Therapeutic communication is the ability or skills of nurses to help clients adapt to stress, overcome psychological problems, and learn how to relate to others (Suryani, 2014). The actual context of the simulation is to improve communication, teamwork, delegation, priority settings, and leadership skills (Bauman *et al*, 2008).

To improve self-confidence and therapeutic communication skills, the most appropriate nursing academy students are using a simulation method that can be standardized patient or role play. Because this simulation method involves all aspects (cognitive, affective and psychomotor) of students in achieving the competencies to be achieved, therapeutic communication skills are a basic competency of nursing students that must be mastered before carrying out the practice of the nursing profession, and appropriate if done on students in the early semester of lecture and this is following the current nursing academy curriculum.

CONCLUSION

The standardized patient method is more effective than the role play method for increasing self-confidence and therapeutic communication skills in nursing students.

SUGGESTIONS

Considering that the main task of the lecturer is to assist students in learning and related to the functions of the lecturer as a designer, instructor, instructor of learning and as a learning evaluator, the lecturer is required to continue to develop themselves in terms of providing facilities that facilitate students to learn, one of them is by using simulation learning strategies. Standardized patient and role play to make it easier to achieve competency standards.

ACKNOWLEDGMENT

Thank you to all respondents and those who have helped in completing this research until the compilation of this manuscript.

DECLARATION OF CONFLICTING INTEREST

There is no conflict of interest in this research.

FUNDING

The funds used in this study came from the researchers' personal funds.

AUTHOR CONTRIBUTION

Bagus Dwi Cahyono: as chairman of the research, coordinate the implementation of research, and compiling scientific article.

Ida Zuhroidah: Conduct research.

Mokhammad Sujarwadi: Conduct research.

ORCID

Bagus Dwi Cahyono

<https://orcid.org/000-0001-5819-6219>

Ida Zuhroidah

<https://orcid.org/0000-0002-4544-0099>

Mokhammad Sujarwadi

<https://orcid.org/0000-0002-3818-0671>

REFERENCES

- Aebersold, M., Tschannen, D., & Bathish, M. (2012). Innovative Simulation Strategies in Education. *Nursing Research and Practice*, 2012. DOI: <http://dx.doi.org/10.1155/2012/765212>.
- Baumann, M., Baumann, C., Le Bihan, E., & Chau, N. (2008). How patients perceive the therapeutic communications skills of their general practitioners, and how that perception affects adherence: use of the TCom-skill GP scale in a specific geographical area. *BMC health services research*, 8(1), 244.
- Blatner, A. (2009). *Role-Playing In Education*. Retrived from <http://www.blatner.com/adam/pdntbk/rpl/ayedu.htm>
- Blum, C. A., Borglund, S., & Parcells, D. (2010). High-fidelity nursing simulation: Impact on student self-confidence and clinical competence. *International Journal of Nursing Education Scholarship*, 7(1) 1-16.
- Bosse, H. M., Nickel, M., Huwendiek, S., Jünger, J., Schultz, J. H., & Nikendei, C. (2010). Peer role-play and standardised patients in communication training: a comparative study on the student perspective on acceptability, realism, and perceived effect. *BMC Medical Education*, 10(1), 27.
- Bowles, N., Mackintosh, C., & Torn, A. (2001). Nurses' communication skills: an evaluation of the impact of solution-focused communication training. *Journal of Advanced Nursing*, 36(3), 347-354.
- Deveugele, M., Derese, A., De Maesschalck, S., Willems, S., Van Driel, M., & De Maeseneer, J. (2005). Teaching communication skills to medical students, a challenge in the curriculum?. *Patient education and counseling*, 58(3), 265-270.
- Fukui, S., Ogawa, K., Ohtsuka, M., & Fukui, N. (2008). A randomized study assessing the efficacy of communication skill training on patients' psychologic distress and coping: nurses' communication with patients just after being diagnosed with cancer. *Cancer: Interdisciplinary International Journal of the American Cancer Society*, 113(6), 1462-1470. DOI: <http://dx.doi.org/10.1002/cncr.23710>.
- Hermawan, A. H. (2010). *Persepsi Pasien Tentang Pelaksanaan Komunikasi Terapeutik Perawat Dalam Asuhan Keperawatan Pada Pasien di Unit Gawat Darurat RS. Mardi Rahayu Kudus November 2009* (Doctoral dissertation, Universitas Diponegoro).
- Hober, C. L. (2012). *Student perceptions of the observer role play experiences in the implementation of a high-fidelity patient simulation in bachelor's degree nursing programs* (Doctoral dissertation, University of Kansas).
- Joyce, B. R., & Weil, M. (2000). *Role Playing: Studying Social Behavior and Values. Models of Teaching. Allyn and Bacon*.
- Jarvis, L., Odell, K., & Troiano, M. (2002). Role-playing as a teaching strategy. *Strategies for application and presentation, staff development and presentation*.
- Lane, C., & Rollnick, S. (2007). The use of simulated patients and role-play in communication skills training: a review of the literature to August 2005. *Patient education and counseling*, 67(1-2), 13-20.
- Lane, C., Hood, K., & Rollnick, S. (2008). Teaching motivational interviewing: using role play is as effective as using simulated patients. *Medical Education*, 42(6), 637-644.
- Mullan, B. A., & Kothe, E. J. (2010). Evaluating a nursing communication skills training course: The relationships between self-rated ability, satisfaction, and actual performance. *Nurse education in practice*, 10(6), 374-378.

- Mäkelä, E., Koistinen, S., Siukola, M., & Turunen, S. (2005). The process model of role playing. *Dissecting LARP*, 205-236.
- Murti, B. (2006). Desain dan ukuran sampel untuk penelitian kuantitatif dan kualitatif di bidang kesehatan. *Yogyakarta: Gadjah Mada University Press*, 67, 113-3.
- Notoatmodjo, S., (2010). *Promosi Kesehatan Teori dan Aplikasi*. Jakarta: Rineka Cipta.
- Neven, R. S. (2010). *Core principles of assessment and therapeutic communication with children, parents and families: Towards the promotion of child and family wellbeing*. Routledge.
- Potter, P.A. & Perry, A.G. (2010). *Fundamental Keperawatan*. Singapore: Elsevier.
- Potter, P.A., Perry, A.G., Stockert P., Hall A. (2014). *Essentials for Nursing Practice. 8th Ed*. St. Louis, Missouri: Mosby Elsevier.
- Rauen, C. A. (2004). Simulation as a teaching strategy for nursing education and orientation in cardiac surgery. *Critical care nurse*, 24(3), 46-51.
- Reading S, Webster B. (2013). *Achieving Competencies for Nursing Practice: A Handbook for Student Nurses*. England: Open University Press.
- Sanford, P.G. (2010). Simulation in Nursing Education: A Review of the Research. *The Qualitative Report*, 15(4), 1006-1011.
- Sleeper, J. A., & Thompson, C. (2008). The use of hi fidelity simulation to enhance nursing students' therapeutic communication skills. *International Journal of Nursing Education Scholarship*, 5(1), 1-12.
- Suikkala, A., Leino-Kilpi, H., & Katajisto, J. (2008). Nursing student-patient relationships: a descriptive study of students' and patients' views. *International Journal of Nursing Education Scholarship*, 5(1), 1-12.
- Supranto, J. (2000). *Teknik Sampling untuk Survei dan Eksperimen*. Jakarta: Rineka Cipta.
- Suryani. (2014). *Komunikasi Terapeutik Teori dan Praktek Edisi 2*. Jakarta: EGC.
- Stuart, G.W., Sundeen, S.J., (2013). *Buku Saku Keperawatan Jiwa Edisi 5*. Jakarta: EGC.
- Üstün, B. (2006). Communication skills training as part of a problem-based learning curriculum. *Journal of Nursing Education*, 45(10).
- Videbeck, S. L. (2008). *Buku Ajar Keperawatan Jiwa*. Jakarta: EGC.
- Von Gunten, C. F., Ferris, F. D., & Emanuel, L. L. (2000). Ensuring competency in end-of-life care: communication and relational skills. *Jama*, 284(23), 3051-3057.
- Waldner, M. H., & Olson, J. K. (2007). Taking the patient to the classroom: Applying theoretical frameworks to simulation in nursing education. *International Journal of Nursing Education Scholarship*, 4(1). DOI: <http://dx.doi.org/10.2202/1548-923X.1317>
- Yedidia, M. J., Gillespie, C. C., Kachur, E., Schwartz, M. D., Ockene, J., Chepaitis, A. E., ... & Lipkin Jr, M. (2003). Effect of communications training on medical student performance. *Jama*, 290(9), 1157-1165.

Cite this article as: Cahyono, B. D., Zuhroidah, I., Sujarwadi, M. (2020). Effectiveness of standardized patients and role play simulation methods in improving self-confidence and therapeutic communication skills of nursing students. *Nurse and Health: Jurnal Keperawatan*, 9(1), 9-15. <https://doi.org/10.36720/nhjk.v9i1.128>