

14. RISK FACTORS FOR THE ANEMIA IN PREGNANT WOMEN: A LITERATURE REVIEW

By Nurnaningsih Nurnaningsih

Review Article: Systematic Review, Meta-Analysis, Integrative Review, Scoping Review

23
RISK FACTORS FOR THE ANEMIA IN PREGNANT WOMEN: A LITERATURE REVIEW

Nurnaningsih^{1*}, Mardiana Ahmad¹, Isharyah Sunarno², Nur Aliya Arsyad¹

11
¹ Midwifery Study Program, Graduate School Hasanuddin University, Makassar, Indonesia

² Department of Obgyn and Gynecology, Faculty of Medicine Hasanuddin University, Makassar, Indonesia

***Correspondence:**

24 **nurnaningsih**
Midwifery Study Program, Graduate School Hasanuddin University, Makassar, Indonesia
Graduate School, **35** Hasanuddin University Makassar, Jln. Perintis Kemerdekaan KM.10 Makassar-90245
Email:
nurnaningsih20p@student.unhas.ac.id

Article Info:

Received: August 17, 2021

Revised: June 21, 2022

Accepted: June 26, 2022

DOI:

<https://doi.org/10.36720/nhjk.v11i1.305>

33
Abstract

Background: Anemia in pregnancy is still a global public health problem and is the largest contributor to morbidity and mortality. Some studies report that risk factors for anemia vary from place to place, so more in-person observations are needed regarding anemia risk factors.

Objectives: This study aimed to determine the risk factors of anemia in pregnant women. Methods of using literature review studies.

Design: This study design is a literature review to search and review article from database and the theory which is descriptive.

Data Sources: Search for scientific articles using the Mendeley application by entering the keyword "Risk Factors, Anemia, Pregnancy" with Published year 2018-2021, English and Indonesian, full text, National journal has ISSN and open access.

Review Methods: The literature review method using narrative review based on inclusion criteria, namely articles discussing anemia in pregnant women, published in 2015-2021, International and National Publications, having ISSN National Journals, Articles using English, original articles, full text and open access. While the Exception Criteria for Articles other than English, type of research literature review and RCT, articles are then collected and a journal summary is made containing the name of the researcher, the year the journal was published, the title of the study, the method and a summary of the results or findings. The summary of research journals is entered into a table according to the format, and 10 articles are eligible.

43
Results: From the 10 selected articles, it can be concluded that the risk factors for anemia, especially in developing countries, are multifactorial, including micro-iron deficiency, folate, and vitamins A and B12 as well as anemia due to parasitic infections such as malaria and hookworms or chronic infections such as TB, HIV, parasites, intestines, infection of working mothers (farmers), contaminated drinking water sources, consumption of coffee/tea and diet. Meanwhile, other factors in Indonesia are unfavorable geographical location, level of education, lack of public awareness of health, socio-economic situation, parity (number of children), and chronic energy shortages (KEK).

17
Conclusion: Anemia is still a global health problem and still has to be addressed, especially in developing countries. Therefore, efforts are needed to better understand the main causes of anemia, including iron deficiency as well as other nutritional deficiencies, diseases, and Hb disorders related to anemia so that appropriate action can be given. Biochemical measurement of micronutrients (especially iron and Vitamin A), inflammatory signs in addition to hematological index when assessing anemia clinically are also urgently needed

Keywords: Risk Factors, Anemia, Pregnancy.

19 INTRODUCTION

Anemia in pregnancy is still a public health problem globally and is the largest contributor to morbidity and mortality (A. et al., 2005; Black et al., 2013; Brabin et al., 2001; Haider & Bhutta, 2017; Keats et al., 2019; Means, 2020; Stephen et al., 2018).

The World Health Organization (WHO) reports about 32.4 million pregnant women suffer from anemia worldwide, with the highest prevalence in Africa (44.6%) followed by Asia with a prevalence of 39.3% (World Health Organization, 2015a).

Anemia is a decrease in the number and size of red blood cells (hemoglobin / Hb concentration) below the set limit value, resulting in a lack of ability for blood to transport oxygen throughout the body (L.T. et al., 2017; Osman et al., 2020; Stephen et al., 2018; WHO, 2006, 2012). About 20% of maternal deaths are caused by anemia and most occur in developing countries. (Black et al., 2013; Kefiyalew et al., 2014; Osman et al., 2020; Stephen et al., 2018). Gestational anemia significantly affected 32.4 million (38.2%) pregnant women. This health problem is most severe in Southeast Asia (48.7%) and Africa (46.3%) (Osman et al., 2020; Stevens et al., 2013; WHO, 2006).

Basic Health Research (2013) reported that the incidence of anemia in pregnant women in Indonesia amounted to 37.1%. (Badan Penelitian dan Pengembangan Kesehatan, 2013) Increased to 48.9% in 2018 (RISKESDAS, 2018).

While the incidence of anemia in Biak Numfor Regency in 2018 was 81% of 3,087 pregnant women, and in 2019 amounted to 79 (%) of 3,063 pregnant women (*Profil Dinas Kesehatan Kab. Biak Numfor*).

It is reported that 40% of the most common cases of anemia are caused by iron

deficiency or about 1.24 billion people worldwide are affected by iron deficiency anemia, but this causative factor is different in each community group and each region (de Leeuw et al., 1966; Means, 2020; Osman et al., 2020; WHO, 2006).

Anemia that occurs in pregnant women has a devastating impact on health, social, and economic status, leading to an increased risk of lack of physical activity, increasing the risk of premature birth, low birth weight, intrauterine fetal death, neonatal death, maternal death, and consequently infant mortality, especially in mothers with severe anemia (Ayano, 2018; Ezzati et al., 2004; Figueiredo et al., 2018; Gedefaw et al., 2015; Haider et al., 2013; Osman et al., 2020; Rahman et al., 2016; Sharma et al., 2020; Zhang et al., 2009).

The study findings report that the risk factors for anemia vary from place to place so more deep observation is needed related to the risk factor of anemia (Osman et al., 2020; WHO, 2006).

Based on the above problems, the research wants to examine more deeply related risk factors for anemia in pregnant women by using literature studies.

METHODS

Design

The design of this research is Literature Review or literature review. The nature of this study is descriptive analysis, which is the regular parsing of data that has been obtained, then given understanding and explanation in order to be well understood by readers.

Search Methods

Literature review is compiled through the search of research articles that have been published both nationally and internationally.

The population and sample were pregnant women with anemia.

Search for scientific articles using the Mendeley application by entering the keyword "Risk Factors, Anemia, Pregnancy" with Published year 2018-2021. Articles selected based on:

a. Inclusion Criteria

1. Article discussing anemia in pregnant women;
2. Published year 2018-2021;
3. International and National Publications;
4. Have an ISSN for National Journal;
5. Articles using English; and
6. Original articles, full text and open access.

b. Exclusion Criteria

1. Articles other than English; and
2. Types of review literature research and RCT.

Search Outcome

From the search obtained 4,124 articles after filtering the year that is 2018-2021 obtained 990 articles, then selecting the category of journals, duplication, open access, research design taken is mix methods study, cross sectional study, correlation analysis, and qualitative study obtained 90 articles.

The final process is to conduct journal selection based on inclusion criteria obtained by 10 journals that are eligible for review. Article Search Strategy can be seen in figure 1.

Quality Appraisal

This literature review is synthesized using narrative methods by grouping similar extraction data according to the results measured to answer the purpose. Research journals that fit the inclusion criteria are then collected and made journal summaries including the name of the researcher, the year of publication of the journal, the title of the study, methods and summary of results or findings (Fadlallah et al., 2019; Pesut et al., 2020).

The summary of the research journal is entered into the table in accordance with the

format mentioned above. To further clarify abstract analysis and full text the journal is read and observed. The journal summary is then conducted an analysis of the content contained in the purpose of the research and the results / findings of the study.

Analysis of the contents of the journal, then coded into the contents of the journal reviewed based on the outline or core of the study is done by parsing in a sentence then if it has been collected then searched for similarities and differences in each study and then discussed to draw conclusions.




Figure 1. Article Search Strategy.

Data Abstraction

Researchers accompanied by two supervisors simultaneously review and read the full article, then discuss to agree on the decision of the article that is adjusted to the criteria of inclusion.

Data Analysis/ Synthesis

After the researchers conducted filtering such as duplication, year of publication, the type of research found ten articles were selected based on inclusion criteria that discussed risk factors for anemia in pregnant women. The filtering process can be seen in Figure 1.

RESULTS

From the search for articles using the Mendeley application, results can be seen on table 1 extraction result of research.

Table 1. Extraction of Research Results.

Title	Author/Year	Type of Research and Sample	Data Analysis	Result
Risk factors of anemia among pregnant women attending antenatal care in health facilities of Eastern Zone of Tigray, Ethiopia	(Berhe et al., 2019)	Control case studies. Sample 600 pregnant women in 2017/2018.	Bivariate and multivariate logistic regression analysis	Risk factors for anemia during pregnancy are significantly intestinal parasites, occupation (farmers), unprotected (polluted) drinking sources, drinking coffee with or immediately after daily meals and diet
Risk Factors for Anemia in Pregnant Women (Study In The Working Area of Slawi Kab. Tegal Health Center)	(Fatkhayah, 2018)	Observational research with case control design. Sample 60 pregnant women divided into 2 groups.	Chi square	There is a positive association of parity on the incidence of anemia during pregnancy.
Prevalence and risk factors of anemia among ever-married women in Jordan	(Arabyat et al., 2019)	Cross sectional. Representative sample of married women aged 15-49 years tested for hemoglobin levels in JPFHS 2012.	Univariate and multivariate logistic regression	Knowledge is related to the incidence of anemia
Spatial distribution and determinant factors of anaemia among women of reproductive age in Ethiopia: a multilevel and spatial analysis	(Kibret et al., 2019)	Observational. 645 cluster of enumeration area (EA).	Multivariable logistic regression model	Factors of risk for anemia are demographics, low education, low income, women living in rural areas.
Anemia in pregnant women participants of the First 1000 Days of Life Program in Agats, Asmat, Papua: Prevalence and analysis of risk factors	(Astari et al., 2018)	Descriptive observational. Sample 230 pregnant women.	Univariate, bivariate, and multivariate.	Risk factors for anemia in pregnant women include the geographic location of Asmat, low level of public education, lack of health workers at health care facilities, lack of public attention to health, low socioeconomic conditions, and nutritional status of pregnant women
Prevalence and Determinant of The Incidence of Anemia pregnant women	(Harna et al., 2020)	Cross sectional. Sample 94 pregnant women.	Descriptive and presented in the form of frequency and cross tabulation	Factors that affect the incidence of anemia in pregnant women are gestational age, KE and parity.
Anaemia in Pregnancy: Prevalence, Risk Factors, and Adverse Perinatal Outcomes in Northern Tanzania	(Stephen et al., 2018)	Cohort study. Sample 539 pregnant women.	Proportions were used for categorical variables and mean or median with respective measures of dispersion for numerical variables. Odds Ratio (OR) with 95%	The mother's low level education is a factor associated with anemia during pregnancy.

			Confidence Interval		
4	Prevalence, risk factors and associated adverse pregnancy outcomes of anaemia in Chinese pregnant women: a multicenter retrospective study	(Lin et al., 2018)	Retrospective study. Sample 44,002 pregnant women.	69) Bivariate and multivariate logistic regression analyses	Anemia is significantly associated with maternal age of 35, family income, rural living BMI.
12	Prevalence of anaemia, risk of haemoglobin concentration and risk factors during the three trimesters of pregnancy	(Arija, 2018)	Longitudinal study. Sample 11,259 women whose pregnancies.	Chi-squared test	The risk factor of anemia is too young and old.
21	Knowledge and Attitude of Pregnant Women in Rural Tanzania on Prevention of Anaemia	(Margwe & Lupindu, 2018)	Cross-sectional. Sample of 354 pregnant women	Fisher's exact test, multivariable logistic regression	Knowledge and attitude related to anemia

DISCUSSION

Anemia is a worldwide public health problem, particularly in developing countries that can affect human health, development, social and economic health (Black et al., 2013; Kassebaum et al., 2016; Mbule et al., 2013; Stephen et al., 2018; Stevens et al., 2013; Vaz-Tostes et al., 2016; O, 2011; WHO and UNICEF, 2004). The goal of the study was to identify risk factors for anemia by considering as many factors as possible.

The results of the journal review identified several risk factors of anemia namely micronutrient deficiency of iron, folate, and vitamins A and B12 and anemia due to parasitic infections such as malaria and hookworms or chronic infections such as TB, HIV, intestinal parasites, maternal work (farmers), contaminated drinking water sources, drinking coffee/tea and diet are major risk factors for anemia in developing countries (Berhe et al., 2019; Foote et al., 2013; Hoffmann et al., 2015; Metz, 2008; Ononge et al., 2014; Stephen et al., 2018; WHO, 2005; Zhang et al., 2009; Hirani and Kamaliani, 2013; Stephen et al., 2018; Dogra, 2020).

While in Indonesia the risk factor of anemia is the unfavorable geographical location, education level, lack of public attention to health, socio-economic circumstances, parity (number of children), adequacy of iron consumption, and KEK status

(Amallia et al., 2017; Astari et al., 2018; Fatkhayah, 2018; Hama et al., 2020; Irwanti et al., 2019; Pasivati & Hatma, 2019; Ristica, 2013; Satriani et al., 2019; Tanziha et al., 2016). In addition to factors above culture is a contributing factor to the occurrence of anemia (Afiyah Sri Harnany, 2006; Contento, 2008).

Several vitamins play a role in iron metabolism, including riboflavin (B2), pyridoxine (B6), cobalamin (B12), and folate. Deficiency of this nutrient has been linked to the incidence of anemia, but the extent of this vitamin's role in anemia is not yet clearly known (Arredondo et al., 2006).

Deficiency of vitamin B12 (cobalamin) and folate can cause macrocytic anemia that can affect DNA synthesis and cell division in the bone marrow (megaloblastic changes), such as hypersegmented neutrophils (Arredondo et al., 2006; Chaparro & Suchdev, 2019; Fishman et al., 2000; Hacibekiroglu et al., 2015; Jafari et al., 2013; Wiering et al., 2016).

In general, iron deficiency is a cause of anemia, although deficiencies in other nutrients and vitamins can also cause anemia, including deficiencies in vitamins A, B12, B6, C, D, and E, folate, riboflavin, copper, and zinc. Some of these vitamins such as vit A, B6, and B12, folic acid, and riboflavin are needed by the body to produce red blood cells normally. While other nutrients, such as vitamins C and E, can protect

red blood cells as anti-oxidants. (Chaparro & Suchdev, 2019; Losken et al., 2005).

Malaria caused by the Plasmodium parasite can cause severe anemia, in addition to other complications including death. (Baird, 2007; Barcus et al., 2007; M. et al., 2007; Sohail et al., 2015; World Health Organization, 2016).

P. falciparum malaria during pregnancy is the leading cause of maternal and fetal morbidity and mortality. Although *P. vivax* infection receives less attention from *P. falciparum* infection, both play an important role in causing anemia in the mother and cause BBLR type of infection to always coexist. (Duffy et al., 2001; Hamer et al., 2009; Sohail et al., 2015).

Parasites require iron for growth, and malaria significantly disrupts iron metabolism in a variety of ways including through hemolysis, heme release, damaged erythropoiesis, increased iron in macrophages, and decreased iron absorption. (Sohail et al., 2015; Spottiswoode et al., 2014).

Anemia is the most common hematological disorder among patients infected with HIV. The disease is usually characterized as normochromic and normocytic anemia with low reticulocyte counts, normal iron storage, and impaired EPO responses.

The prevalence of anemia in HIV-positive people increases along with the development of the disease caused by several factors either indirectly or directly related to the virus. HIV infection causes chronic disease acute phase response, increased hepcidin and AI, and altered iron metabolism (Minchella et al., 2015; Redig & Berliner, 2013; Sohail et al., 2015).

Intestinal parasites (hookworms, *Ascaris lumbricoides*, *Trichuris trichiuria*) are a risk factor for anemia in pregnancy, this is in line with research conducted in Vietnam (Kramer & Kakuma, 2012), Nigeria (Idowu et al., 2005), Benin (Ouédraogo et al., 2012), Ethiopia (Haidar & Pobocik, 2009), Gilgel Gibe area (Getachew et al., 2012), Southeastern Ethiopia (Kefiyalew et al., 2014), Southern Ethiopia

(Lebso et al., 2017), and in Northwestern Ethiopia (Asrie, 2017).

Intestinal parasites cause blood loss, so the mother and fetus are at high risk of anemia. This is because worms in the intestine can cause intestinal necrosis and blood loss due to attachment to the intestinal mucosa (Brooker et al., 2008; World Health Organization, 2015).

Pregnant women who work as farmers are at risk of anemia, which is associated with a lack of information about balanced nutritional intake during pregnancy, economic factors that impact on not being able to access and utilize health services. (EDHS, 2016; Kassa et al., 2017; Xu et al., 2016).

Sources of water that are not clean (polluted), can cause various infections that affect the reduction of iron and other micronutrients and can cause anemia (Berhe et al., 2019).

Consuming coffee/tea after meals immediately after meals, anemia can affect the presence of anti-nutritional factors such as tannins and caffeine, which are found in tea and coffee, respectively (Baig-Ansari et al., 2008; Obse et al., 2013; Taylor & Meyers, 2012).

Tea and coffee can decrease iron absorption by 60% and 50% respectively (Allison, 2000; Anderson & Fitzgerald, 2010). The effect of tea on the absorption of non-hem iron is thought to stem from the formation of an insoluble tannate iron complex (Obse et al., 2013).

In some studies, the level of education has been reported to have an effect on the reduced risk of anemia. Educated pregnant women have a better income so that it affects the food to be consumed so that they do not experience nutritional anemia (Tanzania Bureau of Statistics, 2010).

It was further reported in Ethiopia that the prevalence of anemia was higher in uneducated pregnant women (Melku et al., 2014). Secondary and higher education levels are also associated with maternal compliance in terms of exclusive breastfeeding, ANC visits in accordance with the recommendations given (Tanzania Bureau of Statistics, 2010).

Income levels also affect the incidence of anemia, this is because low income leads to an inability to consume nutritious foods. A study in Ethiopia showed that women with low incomes are more at risk of suffering from anemia than women with higher incomes (Bekele et al., 2016; Gedefaw et al., 2015; Stephen et al., 2018).

Parity is a factor in the occurrence of anemia in pregnant women, this is associated with the theory that a mother who often gives birth has a risk of anemia in subsequent pregnancies if she does not pay attention to nutritional needs, because during pregnancy nutrients will share for the mother and the fetus they contain. The more often a woman gives birth, the greater the risk of blood loss and has an impact on decreasing Hb levels. Each time a woman gives birth, the amount of iron lost is estimated at 250mg (Pasmawati & Hatma, 2019).

Adequacy of iron consumption related to the occurrence of anemia is due to the need for iron in pregnancy increased, especially in the final trimester, in the process of maturation of iron red blood cells taken from plasma transferrin which is the iron reserve in serum. (Pasmawati & Hatma, 2019).

Pregnant women with an arm circumference (LILA) of less than 23 cm are at risk of anemia. (Milman et al., 2000). The upper arm is the measure used for nutritional status assessment, with the normal limit being 23 cm for adult women, while the lesser arm circumference is also associated with undernutrition status. This is in line with the results of studies reported in eastern Ethiopia and western Ethiopia, Kenya, Nepal, and India (Alene & Mohamed Dohe, 2015; Mondal et al., 2006; Osman et al., 2020). With the similarity of reported research results, malnourished pregnant women are at higher risk of anemia.

Culture is one of the factors of anemia. Culture is the experience of people from the moment they are born, culture is also shared knowledge and shared meaning, where the meaning indicates the complexity of belief or knowledge and the connection of values or

feelings with beliefs (Contento, 2008). The results of health ethnographic research in 2012 on 12 ethnicities in Indonesia showed that maternal and child health problems are still very concerning.

In Mexico, pregnant women and after childbirth are prohibited from eating foods that are "cold". In Indonesia alone pregnant women and after childbirth are prohibited from eating eggs, meat, shrimp, sea fish and catfish, conch, lembayung leaves, bitter melon, pineapple, brown sugar, and foods fried with oil (Hamany, 2006).

So that by knowing the risk factors for anemia, health workers can implement policies that focus more on the risk group and perform treatment for insufficient iron and provide appropriate iron supplements. With proper treatment is expected to reduce the number of deaths and the number of pains caused by anemia.

CONCLUSION

Anemia is still a global health problem and still has to be addressed especially in developing countries. So, efforts are needed to better understand the main causes of anemia including iron deficiency as well as other nutritional deficiencies, diseases, and Hb disorders associated with anemia so that appropriate measures can be given. Biochemical measurements of micronutrients (especially iron and Vitamin A), signs of inflammation, in addition to the hematological index when assessing anemia clinically are also urgently needed.

ACKNOWLEDGMENT

Thank you to the supervisor who has taken the time to direct in the preparation of literature studies.

DECLARATION OF CONFLICTING INTEREST

This comprehensive summary or thematic review is independent writing, so there is no conflict of interest in the writing.

FUNDING

This systematic review is writing independently, not funded or get funding from any party.

AUTHOR CONTRIBUTION

Nurnaningsih: Designed the study, collected and analyzed articles, and contributed to the completion of a systematic review.

Mardiana Ahmad: Contribution as a supervisor guide and discuss the final results of the review literature manuscript.

Isharyah Sunarno: Contribution as a supervisor involved in planning and supervision in the completion of the literature review.

Nur Aliya Arsyad: Contributed to the preparation of the final manuscript.

ORCID

Yohana Rita Bitbit

None.

Mardiana Ahmad

<https://orcid.org/0000-0002-0798-0457>

Isharyah Sunarno

None.

Nur Aliya Arsyad

None.

REFERENCES

- A., L., D., F., M., K., M., M., & E., S. (2005). Maternal anemia during pregnancy is an independent risk factor for low birthweight and preterm delivery. *European Journal of Obstetrics and Gynecology and Reproductive Biology*.
- Afiyah Sri Harnany. (2006). *PENGARUH TABU MAKANAN, TINGKAT KECUKUPAN GIZI, KONSUMSI TABLET BESI, DAN TEH TERHADAP KADAR HEMOGLOBIN PADA IBU HAMIL DI KOTA PEKALONGAN*

TAHUN 2006. UNIVERSITAS DIPONEGORO.

- Alene, K. A., & Mohamed Dohe, A. (2015). Prevalence of Anemia and Associated Factors among Pregnant Women in an Urban Area of Eastern Ethiopia. *Anemia*. <https://doi.org/10.1155/2014/561567>
- Allen, L. H. (2000). Anemia and iron deficiency: Effects on pregnancy outcome. *American Journal of Clinical Nutrition*. <https://doi.org/10.1093/ajcn/71.5.1280s>
- Amallia, S., Afriyani, R., & Utami, S. P. (2017). Faktor Risiko Kejadian Anemia pada Ibu Hamil di Rumah Sakit BARI Palembang. *Jurnal Kesehatan*. <https://doi.org/10.26630/jk.v8i3.639>
- Anderson, J., & Fitzgerald, C. (2010). Iron : An Essential Nutrient. *Food and Nutrition Series/Health*.
- Arabyat, R., Arabyat, G., & Al-Taani, G. (2019). Prevalence and risk factors of anaemia among ever-married women in Jordan. *Eastern Mediterranean Health Journal*. <https://doi.org/10.26719/emhj.18.074>
- Arija, V. (2018). PREVALENCE OF ANAEMIA, RISK OF HAEMOCONCENTRATION AND RISK FACTORS DURING THE THREE TRIMESTERS OF PREGNANCY. *Nutrición Hospitalaria*. <https://doi.org/10.20960/nh.1045>
- Arredondo, M., Martínez, R., Núñez, M. T., Ruz, M., & Olivares, M. (2006). Inhibition of iron and copper uptake by iron, copper and zinc. *Biological Research*. <https://doi.org/10.4067/S0716-97602006000100011>
- Asrie, F. (2017). Prevalence of anemia and its associated factors among pregnant women receiving antenatal care at Aymiba Health Center, northwest Ethiopia. *Journal of Blood Medicine*. <https://doi.org/10.2147/JBM.S134932>
- Astari, P., Rinonce, H. T., Kasim, F., Pudjohartono, M. F., Debora, J., & Winata, M. G. (2018). Anemia pada ibu

- hamil peserta Program 1000 Hari Pertama Kehidupan di Agats, Asmat, Papua: Prevalensi dan analisis faktor risiko. *Journal of Community Empowerment for Health*.
<https://doi.org/10.22146/jcoemph.39261>
- Ayano, B. (2018). Assessment of Prevalence and Risk Factors for Anemia Among Pregnant Mothers Attending Anc Clinic at Adama Hospital Medical Collage, Adama, Ethiopia, 2017. *Journal of Gynecology and Obstetrics*.
<https://doi.org/10.11648/j.jgo.20180603.11>
- Badan Penelitian dan Pengembangan Kesehatan. (2013). Riset Kesehatan Dasar 2013. *Riset Kesehatan Dasar 2013*.
- Baig-Ansari, N., Badruddin, S. H., Karmaliani, R., Harris, H., Jehan, I., Pasha, O., Moss, N., McClure, E. M., & Goldenberg, R. L. (2008). Anemia prevalence and risk factors in pregnant women in an urban area of Pakistan. *Food and Nutrition Bulletin*.
<https://doi.org/10.1177/156482650802900207>
- Baird, J. K. (2007). Neglect of Plasmodium vivax malaria. In *Trends in Parasitology*.
<https://doi.org/10.1016/j.pt.2007.08.011>
- Barcus, M. J., Basri, H., Picarima, H., Manyakori, C., Sekartuti, Elyazar, I., Bangs, M. J., Maguire, J. D., & Baird, J. K. (2007). Demographic risk factors for severe and fatal vivax and falciparum malaria among hospital admissions in northeastern Indonesian Papua. *American Journal of Tropical Medicine and Hygiene*.
<https://doi.org/10.4269/ajtmh.2007.77.984>
- Bekele, A., Tilahun, M., & Mekuria, A. (2016). Prevalence of Anemia and Its Associated Factors among Pregnant Women Attending Antenatal Care in Health Institutions of Arba Minch Town, Gamo Gofa Zone, Ethiopia: A Cross-Sectional Study. *Anemia*.
<https://doi.org/10.1155/2016/1073192>
- Berhe, K., Fseha, B., Gebremariam, G., Teame, H., Etsay, N., Welu, G., & Tsegay, T. (2019). Risk factors of anemia among pregnant women attending antenatal care in health facilities of eastern zone of tigray, Ethiopia, case-control study, 2017/18. *Pan African Medical Journal*.
<https://doi.org/10.11604/pamj.2019.34.121.15999>
- Black, R. E., Victora, C. G., Walker, S. P., Bhutta, Z. A., Christian, P., De Onis, M., Ezzati, M., Grantham-Mcgregor, S., Katz, J., Martorell, R., & Uauy, R. (2013). Maternal and child undernutrition and overweight in low-income and middle-income countries. In *The Lancet*.
[https://doi.org/10.1016/S0140-6736\(13\)60937-X](https://doi.org/10.1016/S0140-6736(13)60937-X)
- Brabin, B. J., Hakimi, M., & Pelletier, D. (2001). An analysis of anemia and pregnancy-related maternal mortality. *Journal of Nutrition*.
<https://doi.org/10.1093/jn/131.2.604s>
- Brooker, S., Hotez, P. J., & Bundy, D. A. P. (2008). Hookworm-related anaemia among pregnant women: A systematic review. In *PLoS Neglected Tropical Diseases*.
<https://doi.org/10.1371/journal.pntd.0000291>
- Chaparro, C. M., & Suchdev, P. S. (2019). Anemia epidemiology, pathophysiology, and etiology in low- and middle-income countries. In *Annals of the New York Academy of Sciences*.
<https://doi.org/10.1111/nyas.14092>
- Contento, I. R. (2008). Nutrition education: Linking research, theory, and practice. In *Asia Pacific Journal of Clinical Nutrition*.
<https://doi.org/10.1097/FCH.0b013e3182385c56>
- de Leeuw, N. K. M., Lowenstein, L., & Hsieh, Y. S. (1966). Iron deficiency and hydremia in normal pregnancy. *Medicine (United States)*.
<https://doi.org/10.1097/00005792-196607000-00002>
- Duffy, P. E., Craig, A. G., & Baruch, D. I.

- (2001). Variant proteins on the surface of malaria-infected erythrocytes - Developing vaccines. *Trends in Parasitology*.
[https://doi.org/10.1016/S1471-4922\(01\)02022-0](https://doi.org/10.1016/S1471-4922(01)02022-0)
- EDHS. (2016). Ethiopia 2016 Demographic and Health Survey. *Ethiopia Demographic and Health Survey*.
- Ezzati, M., Lopez, A. D., Rodgers, A., & Murray, C. J. (2004). Comparative Quantification of Health Risks Global and Regional Burden of Disease. In *Comparative Quantification of Health Risks Global and Regional Burden of Disease*.
- Fadlallah, R., El-Jardali, F., Nomier, M., Hemadi, N., Arif, K., Langlois, E. V., & Akl, E. A. (2019). Using narratives to impact health policy-making: A systematic review. In *Health Research Policy and Systems*.
<https://doi.org/10.1186/s12961-019-0423-4>
- Fatkhiyah, N. (2018). FAKTOR RISIKO KEJADIAN ANEMIA PADA IBU HAMIL (STUDI DI WILAYAH KERJA PUSKESMAS SLAWI KAB. TEGAL). *Indonesia Jurnal Kebidanan*.
<https://doi.org/10.26751/ijb.v2i2.561>
- Figueiredo, A. C. M. G., Gomes-Filho, I. S., Silva, R. B., Pereira, P. P. S., Da Mata, F. A. F., Lyrio, A. O., Souza, E. S., Cruz, S. S., & Pereira, M. G. (2018). Maternal anemia and low birth weight: A systematic review and meta-analysis. In *Nutrients*.
<https://doi.org/10.3390/nu10050601>
- Fishman, S. M., Christian, P., & West, J. (2000). The role of vitamins in the prevention and control anaemia. *Public Health Nutrition*.
<https://doi.org/10.1017/s1368980000000173>
- Foote, E. M., Sullivan, K. M., Ruth, L. J., Oremo, J., Sadumah, I., Williams, T. N., & Suchdev, P. S. (2013). Determinants of anemia among preschool children in rural, western Kenya. *American Journal of Tropical Medicine and Hygiene*.
<https://doi.org/10.4269/ajtmh.12-0560>
- Gedefaw, L., Ayele, A., Asres, Y., & Mossie, A. (2015). Anemia and Associated Factors Among Pregnant Women Attending Antenatal Care Clinic in Wolayita Sodo Town, Southern Ethiopia. *Ethiopian Journal of Health Sciences*.
<https://doi.org/10.4314/ejhs.v25i2.8>
- Getachew, M., Yewhalaw, D., Tafess, K., Getachew, Y., & Zeynudin, A. (2012). Anaemia and associated risk factors among pregnant women in Gilgel Gibe dam area, Southwest Ethiopia. *Parasites and Vectors*.
<https://doi.org/10.1186/1756-3305-5-296>
- Hacibekiroglu, T., Basturk, A., Akinci, S., Bakanay, S. M., Ulas, T., Guney, T., & Dilek, I. (2015). Evaluation of serum levels of zinc, copper, and Helicobacter pylori IgG and IgA in iron deficiency anemia cases. *European Review for Medical and Pharmacological Sciences*.
- Haider, J. A., & Pobocik, R. S. (2009). Iron deficiency anemia is not a rare problem among women of reproductive ages in Ethiopia: A community based cross sectional study. *BMC Blood Disorders*.
<https://doi.org/10.1186/1471-2326-9-7>
- Haider, B. A., & Bhutta, Z. A. (2017). Multiple-micronutrient supplementation for women during pregnancy. In *Cochrane Database of Systematic Reviews*.
<https://doi.org/10.1002/14651858.CD004905.pub5>
- Haider, B. A., Olofin, I., Wang, M., Spiegelman, D., Ezzati, M., & Fawzi, W. W. (2013). Anaemia, prenatal iron use, and risk of adverse pregnancy outcomes: Systematic review and meta-analysis. In *BMJ (Online)*.
<https://doi.org/10.1136/bmj.f3443>
- Hamer, D. H., Singh, M. P., Wylie, B. J., Yeboah-Antwi, K., Tuchman, J., Desai, M., Udhayakumar, V., Gupta, P., Brooks, M. I., Shukla, M. M., Awasthy, K., Sabin, L., MacLeod, W. B., Dash, A. P., & Singh, N. (2009). Burden of malaria in pregnancy

- in Jharkhand State, India. *Malaria Journal*. <https://doi.org/10.1186/1475-2875-8-210>
- Harna, Muliani, E. Y., Sa'pang, M., Dewanti, L. P., Muh, A., & Irwan, A. (2020). Prevalensi Dan Determinan Kejadian Anemia Ibu Hamil. *JIK (Jurnal Ilmu Keperawatan)*.
- Harnany, A. S. (2006). *Hubungan Tabu Makanan dan Asupan Zat Gizi dengan Kejadian Anemia pada Ibu Hamil di Kota Pekalongan*. Universitas Diponegoro Semarang.
- Hoffmann, J. J. M. L., Urrechaga, E., & Aguirre, U. (2015). Discriminant indices for distinguishing thalassemia and iron deficiency in patients with microcytic anemia: A meta-analysis. *Clinical Chemistry and Laboratory Medicine*. <https://doi.org/10.1515/cclm-2015-0179>
- Idowu, O. A., Mafiana, C. F., & Dapo, S. (2005). Anaemia in pregnancy: A survey of pregnant women in Abeokuta, Nigeria. *African Health Sciences*. <https://doi.org/10.5555/afhs.2005.5.4.295>
- Irwanti, L., Sulistiyani, S., & Rohmawati, N. (2019). DETERMINAN KEJADIAN ANEMIA GIZI BESI PADA IBU HAMIL KURANG ENERGI KRONIS DI WILAYAH KERJA PUSKESMAS SUMBERJAMBE KABUPATEN JEMBER. *IKESMA*. <https://doi.org/10.19184/ikesma.v15i2.17553>
- Jafari, S. M., Heidari, G., Nabipour, I., Amirinejad, R., Assadi, M., Bargahi, A., Akbarzadeh, S., Tahmasebi, R., & Sanjdideh, Z. (2013). Serum retinol levels are positively correlated with hemoglobin concentrations, independent of iron homeostasis: A population-based study. *Nutrition Research*. <https://doi.org/10.1016/j.nutres.2013.02.004>
- Kassa, G. M., Muche, A. A., Berhe, A. K., & Fekadu, G. A. (2017). Prevalence and determinants of anemia among pregnant women in Ethiopia; a systematic review and meta-analysis. *BMC Hematology*. <https://doi.org/10.1186/s12878-017-0090-z>
- Kassebaum, N. J., Fleming, T. D., Flaxman, A., Phillips, D. E., Steiner, C., Barber, R. M., Hanson, S. W., Moradi-Lakeh, M., Coffeng, L. E., Haagsma, J., Kyu, H. H., Graetz, N., Lim, S. S., Vos, T., Naghavi, M., Murray, C., Yonemoto, N., Singh, J., Jonas, J. B., ... Levy, T. S. (2016). The Global Burden of Anemia. In *Hematology/Oncology Clinics of North America*. <https://doi.org/10.1016/j.hoc.2015.11.002>
- Keats, E. C., Haider, B. A., Tam, E., & Bhutta, Z. A. (2019). Multiple-micronutrient supplementation for women during pregnancy. In *Cochrane Database of Systematic Reviews*. <https://doi.org/10.1002/14651858.CD004905.pub6>
- Kefiyalew, F., Zemene, E., Asres, Y., & Gedefaw, L. (2014). Anemia among pregnant women in Southeast Ethiopia: Prevalence, severity and associated risk factors. *BMC Research Notes*. <https://doi.org/10.1186/1756-0500-7-771>
- Kibret, K. T., Chojenta, C., D'Arcy, E., & Loxton, D. (2019). Spatial distribution and determinant factors of anaemia among women of reproductive age in Ethiopia: A multilevel and spatial analysis. *BMJ Open*. <https://doi.org/10.1136/bmjopen-2018-027276>
- Kramer, M. S., & Kakuma, R. (2012). Optimal duration of exclusive breastfeeding (Review) Optimal duration of exclusive breastfeeding. *Biostatistics*, 8. <https://doi.org/10.1002/14651858.CD003517.pub2>. Copyright
- L.T., N., I., S., B., S.-P., S., P., I., A.-Z., M., R., A.F., J., S., V., Nyflot, L. T., Sandven, I., Stray-Pedersen, B., Pettersen, S., Al-Zirqi, I., Rosenberg, M., Jacobsen, A. F., & Vangen, S. (2017). Risk factors for severe postpartum hemorrhage: A case-control study. *BMC Pregnancy and Childbirth*.
- Lebso, M., Anato, A., & Loha, E. (2017).

- Prevalence of anemia and associated factors among pregnant women in Southern Ethiopia: A community based cross-sectional study. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0188783>
- Lin, L., Wei, Y., Zhu, W., Wang, C., Su, R., Feng, H., & Yang, H. (2018). Prevalence, risk factors and associated adverse pregnancy outcomes of anaemia in Chinese pregnant women: A multicentre retrospective study. *BMC Pregnancy and Childbirth*, 18(1). <https://doi.org/10.1186/s12884-018-1739-8>
- Losken, A., Fishman, I., Denson, D., Moyer, H., & Carlson, G. (2005). An objective evaluation of breast symmetry using 3-dimensional images. *Annals of Plastic Surgery*, 55(6), 571–575.
- M., D., F.O., ter K., F., N., R., M., K., A., B., B., & R.D., N. (2007). Epidemiology and burden of malaria in pregnancy. In *Lancet Infectious Diseases*.
- Margwe, J. A., & Lupindu, A. M. (2018). Knowledge and attitude of pregnant women in rural Tanzania on prevention of anaemia. *African Journal of Reproductive Health*, 22(3). <https://doi.org/10.29063/ajrh2018/v22i3.8>
- Mbule, M. A., Byaruhanga, Y. B., Kabahenda, M., & Lubowa, A. (2013). Determinants of anaemia among pregnant women in rural Uganda. *Rural and Remote Health*. <https://doi.org/10.22605/rrh2259>
- Means, R. T. (2020). Iron deficiency and iron deficiency anemia: Implications and impact in pregnancy, fetal development, and early childhood parameters. In *Nutrients*. <https://doi.org/10.3390/nu12020447>
- Melku, M., Addis, Z., Alem, M., & Enawgaw, B. (2014). Prevalence and predictors of maternal anemia during pregnancy in Gondar, Northwest Ethiopia: An institutional based cross-sectional study. *Anemia*. <https://doi.org/10.1155/2014/108593>
- Metz, J. (2008). A high prevalence of biochemical evidence of vitamin B12 or folate deficiency does not translate into a comparable prevalence of anemia. *Food and Nutrition Bulletin*. <https://doi.org/10.1177/15648265080292s111>
- Milman, N., Byg, K. E., & Agger, A. O. (2000). Hemoglobin and erythrocyte indices during normal pregnancy and postpartum in 206 women with and without iron supplementation. *Acta Obstetrica et Gynecologica Scandinavica*. <https://doi.org/10.1034/j.1600-0412.2000.079002089.x>
- Minchella, P. A., Armitage, A. E., Darboe, B., Jallow, M. W., Drakesmith, H., Jaye, A., Prentice, A. M., & McDermid, J. M. (2015). Elevated hepcidin is part of a complex relation that links mortality with iron homeostasis and anemia in men and women with HIV infection. *Journal of Nutrition*. <https://doi.org/10.3945/jn.114.203158>
- Mondal, B., Tripathy, V., & Gupta, R. (2006). Risk factors of Anemia during pregnancy among the Garo of Meghalaya, India. *J Hum Ecol*, 14(14), 27–32.
- Obse, N., Mossie, A., & Gobena, T. (2013). Magnitude of anemia and associated risk factors among pregnant women attending antenatal care in Shalla Woreda, West Arsi Zone, Oromia Region, Ethiopia. *Ethiopian Journal of Health Sciences*.
- Ononge, S., Campbell, O., & Mirembe, F. (2014). Haemoglobin status and predictors of anaemia among pregnant women in Mpigi, Uganda. *BMC Research Notes*. <https://doi.org/10.1186/1756-0500-7-712>
- Osman, M. O., Nour, T. Y., Bashir, H. M., Roble, A. K., Nur, A. M., & Abdilahi, A. O. (2020). Risk factors for anemia among pregnant women attending the antenatal care unit in selected jigjiga public health facilities, somali region, east ethiopia 2019: Unmatched case-control study. *Journal of Multidisciplinary Healthcare*. <https://doi.org/10.2147/JMDH.S260398>

- Ouédraogo, S., Koura, G. K., Accrombessi, M. M. K., Bodeau-Livinec, F., Massougbodji, A., & Cot, M. (2012). Maternal anemia at first antenatal visit: Prevalence and risk factors in a malaria-endemic area in Benin. *American Journal of Tropical Medicine and Hygiene*. <https://doi.org/10.4269/ajtmh.2012.11-0706>
- Pasmawati, P., & Hatma, R. D. (2019). Determinan Anemia Ibu Hamil Trimester II dan III di Indonesia (Analisis Data Riset Kesehatan Dasar). *Jurnal Kesehatan*. <https://doi.org/10.26630/jk.v10i1.1178>
- Pesut, B., Greig, M., Thorne, S., Storch, J., Burgess, M., Tishelman, C., Chambaere, K., & Janke, R. (2020). Nursing and euthanasia: A narrative review of the nursing ethics literature. In *Nursing Ethics*. <https://doi.org/10.1177/0969733019845127>
- Profil Dinas Kesehatan Kab. Biak Numfor*. (n.d.).
- Rahman, M. M., Abe, S. K., Rahman, M. S., Kanda, M., Narita, S., Bilano, V., Ota, E., Gilmour, S., & Shibuya, K. (2016). Maternal anemia and risk of adverse birth and health outcomes in low- and middle-income countries: Systematic review and meta-analysis. *American Journal of Clinical Nutrition*. <https://doi.org/10.3945/ajcn.115.107896>
- Redig, A. J., & Berliner, N. (2013). Pathogenesis and clinical implications of HIV-related anemia in 2013. In *Hematology / the Education Program of the American Society of Hematology. American Society of Hematology. Education Program*. <https://doi.org/10.1182/asheducation-2013.1.377>
- RISKESDAS. (2018). Riset Kesehatan Dasar 2018. *Kementrian Kesehatan Republik Indonesia*.
- Ristica, O. D. (2013). Faktor Risiko Kejadian Anemia pada Ibu Hamil. *Jurnal Kesehatan Komunitas*. <https://doi.org/10.25311/keskom.vol2.iss2.49>
- Satriani, Veni, H., & Nilawati, A. (2019). Hubungan Faktor Pendidikan Dan Faktor Ekonomi Orang Tua Dengan Kejadian Anemia Pada Remaja Usia 12-18 Tahun Di Kecamatan Tamalatea Kabupaten Jeneponto. *Jurnal JKFT*.
- Sharma, S., Kaur, S. P., & Lata, G. (2020). Anemia in Pregnancy is Still a Public Health Problem: A Single Center Study with Review of Literature. *Indian Journal of Hematology and Blood Transfusion*. <https://doi.org/10.1007/s12288-019-01187-6>
- Sohail, M., Shakeel, S., Kumari, S., Bharti, A., Zahid, F., Anwar, S., Singh, K. P., Islam, M., Sharma, A. K., Lata, S., Ali, V., Adak, T., Das, P., & Raziuddin, M. (2015). Prevalence of malaria infection and risk factors associated with anaemia among pregnant women in semiurban community of Hazaribag, Jharkhand, India. *BioMed Research International*. <https://doi.org/10.1155/2015/740512>
- Spottiswoode, N., Duffy, P. E., & Drakesmith, H. (2014). Iron, anemia and hepcidin in malaria. In *Frontiers in Pharmacology*. <https://doi.org/10.3389/fphar.2014.00125>
- Stephen, G., Mgongo, M., Hussein Hashim, T., Katanga, J., Stray-Pedersen, B., & Msuya, S. E. (2018). Anaemia in Pregnancy: Prevalence, Risk Factors, and Adverse Perinatal Outcomes in Northern Tanzania. *Anemia*. <https://doi.org/10.1155/2018/1846280>
- Stevens, G. A., Finucane, M. M., De-Regil, L. M., Paciorek, C. J., Flaxman, S. R., Branca, F., Peña-Rosas, J. P., Bhutta, Z. A., & Ezzati, M. (2013). Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995-2011: A systematic analysis of population-representative data. *The Lancet Global Health*. [https://doi.org/10.1016/S2214-109X\(13\)70001-9](https://doi.org/10.1016/S2214-109X(13)70001-9)

- Tanzania Bureau of Statistics. (2010). Tanzania Demographic and Health Survey 2010. *National Bureau of Statistics Dar Es Salaam, Tanzania ICF Macro Calverton, Maryland, USA*.
- Tanziha, I., Rizal, M., Damanik, M., Juntra Utama, L., Rosmiati, R., Masyarakat, D. G., & Manusia, F. E. (2016). FAKTOR RISIKO ANEMIA IBU HAMIL DI INDONESIA (Anemia risk factors among pregnant women in Indonesia). In *Jurnal Gizi dan Pangan*.
- Taylor, C. L., & Meyers, L. D. (2012). Advanced Nutrition and Human Metabolism Fifth. *Modern Nutrition in Health and Disease: Eleventh Edition*.
- Vaz-Tostes, M. das G., Verediano, T. A., de Mejia, E. G., & Brunoro Costa, N. M. (2016). Worldwide prevalence of anaemia, WHO Vitamin and Mineral Nutrition Information System, 1993-2005. *Journal of the Science of Food and Agriculture*.
<https://doi.org/10.1039/c1ay05430a>
- WHO. (2005). Worldwide prevalence of anaemia 1993-2005: WHO Global database of Anemia. *WHO Report*.
- WHO. (2006). Neonatal and perinatal mortality: Country region and Global estimates. In *World Health Organisation*.
- WHO. (2011). The global prevalence of anaemia in 2011. *Who*.
- WHO. (2012). *Guideline: Daily Iron and Folic Acid Supplementation in Pregnant Women*.
<http://www.who.int/publications/guidelines/en/>
- WHO and UNICEF. (2004). Focusing on anaemia: towards an integrated approach for effective anaemia control. *Joint Statement by the World Health Organization and the United Nations Children's Fund*.
- Wieringa, F. T., Dahl, M., Chamnan, C., Poirot, E., Kuong, K., Sophonneary, P., Sinuon, M., Greuffeille, V., Hong, R., Berger, J., Dijkhuizen, M. A., & Lailou, A. (2016). The high prevalence of anemia in cambodian children and women cannot be satisfactorily explained by nutritional deficiencies or hemoglobin disorders. *Nutrients*.
<https://doi.org/10.3390/nu8060348>
- World Health Organization. (2015a). *The global prevalence of anaemia in 2011*. WHO.
- World Health Organization. (2015b). WHO Iron deficiency anaemia: assessment, prevention and control. In *WHO*.
- World Health Organization. (2016). *Malaria Fact Sheet*. Malaria Fact Sheet.
- Xu, X., Liu, S., Rao, Y., Shi, Z., Wang, L. L., Sharma, M., & Zhao, Y. (2016). Prevalence and sociodemographic and lifestyle determinants of anemia during pregnancy: A cross-sectional study of pregnant women in China. *International Journal of Environmental Research and Public Health*.
<https://doi.org/10.3390/ijerph13090908>
- Zhang, Q., Ananth, C. V., Rhoads, G. G., & Li, Z. (2009). The Impact of Maternal Anemia on Perinatal Mortality: A Population-based, Prospective Cohort Study in China. *Annals of Epidemiology*.
<https://doi.org/10.1016/j.annepidem.2009.06.002>

Cite this article as: Nurnaningsih, Ahmad, M., Sunarno, I., Arsyad, N.A. (2022). Risk factors for the anemia in pregnant women: A literature review. *Nurse and Health: Jurnal Keperawatan*, 11 (1), 137-150. <https://doi.org/10.36720/nhjk.v11i1.305>

14. RISK FACTORS FOR THE ANEMIA IN PREGNANT WOMEN: A LITERATURE REVIEW

ORIGINALITY REPORT

24%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|---|--|---------------|
| 1 | www.ru.ac.bd
Internet | 61 words — 2% |
| 2 | Indah Nur Imamah. "The Effect of Autogenic Relaxation on Blood Pressure Changes in Patients with Hypertension: Literature Review", STRADA Jurnal Ilmiah Kesehatan, 2020
Crossref | 36 words — 1% |
| 3 | journals.plos.org
Internet | 34 words — 1% |
| 4 | www.wjgnet.com
Internet | 32 words — 1% |
| 5 | Www.dovepress.com
Internet | 26 words — 1% |
| 6 | "Prevalence of Malaria Infection and Risk Factors Associated with Anaemia among Pregnant Women in Semiurban Community of Hazaribag, Jharkhand, India", 'Hindawi Limited', 2015
Internet | 25 words — 1% |
| 7 | Nyasiro S Gibore, Agatha F Ngowi, Mariam J Munyogwa, Mwanaisha M Ali. "Dietary Habits Associated with Anemia Among Pregnant Women Attending | 25 words — 1% |

Antenatal Care Services", Current Developments in Nutrition, 2020

Crossref

8	citeseerx.ist.psu.edu Internet	24 words — 1%
9	informe.ensp.fiocruz.br Internet	22 words — 1%
10	pubmed.ncbi.nlm.nih.gov Internet	22 words — 1%
11	jurnal.upertis.ac.id Internet	20 words — < 1%
12	hdl.handle.net Internet	19 words — < 1%
13	ir.kdu.ac.lk Internet	19 words — < 1%
14	scitepress.org Internet	19 words — < 1%
15	lppm.uhb.ac.id Internet	18 words — < 1%
16	Awoke Kebede, Hadgu Gerensea, Freweyni Amare, Yared Tesfay, Girmay Teklay. "The magnitude of anemia and associated factors among pregnant women attending public institutions of Shire Town, Shire, Tigray, Northern Ethiopia, 2018", BMC Research Notes, 2018 Crossref	17 words — < 1%
17	oaji.net Internet	17 words — < 1%

18	ogma.newcastle.edu.au Internet	17 words — < 1%
19	www.ajol.info Internet	17 words — < 1%
20	www.dovepress.com Internet	16 words — < 1%
21	WWW.Suaire.sua.ac.tz Internet	15 words — < 1%
22	ejournal.urindo.ac.id Internet	15 words — < 1%
23	Mohammadmahdi Karami, Maryam Chaleshgar, Nader Salari, Hakimeh Akbari, Masoud Mohammadi. "Global Prevalence of Anemia in Pregnant Women: A Comprehensive Systematic Review and Meta-Analysis", Maternal and Child Health Journal, 2022 Crossref	14 words — < 1%
24	Musniati, Saidah Syamsuddin, Muhammad Tamar. "Estradiol levels in postpartum blues incidences at Pertiwi Maternal and Child Hospitals in Makassar", Enfermería Clínica, 2020 Crossref	14 words — < 1%
25	www.emro.who.int Internet	14 words — < 1%
26	jurnal.ugm.ac.id Internet	13 words — < 1%
27	udsspace.uds.edu.gh Internet	13 words — < 1%

28 Brhane Berhe, Fitsum Mardu, Haftom Legese, Aderajew Gebrewahd et al. "Prevalence of anemia and associated factors among pregnant women in Adigrat General Hospital, Tigray, northern Ethiopia, 2018", BMC Research Notes, 2019

12 words — < 1%

Crossref

29 Mohamed Omar Osman, Tahir Yousuf Nour, Hodan Mahamed Bashir, Abdurahman Kedir Roble et al. "<p>Risk Factors for Anemia Among Pregnant Women Attending the Antenatal Care Unit in Selected Jigjiga Public Health Facilities, Somali Region, East Ethiopia 2019: Unmatched Case–Control Study</p>", Journal of Multidisciplinary Healthcare, 2020

12 words — < 1%

Crossref

30 apps.who.int

Internet

12 words — < 1%

31 www.mdpi.com

Internet

12 words — < 1%

32 Alemayehu Digssie Gebremariam, Sofonyas Abebaw Tiruneh, Bedilu Abebe Abate, Melaku Tadege Engidaw, Desalegn Tesfa Asnakew. "Adherence to iron with folic acid supplementation and its associated factors among pregnant women attending antenatal care follow up at Debre Tabor General Hospital, Ethiopia, 2017", PLOS ONE, 2019

11 words — < 1%

Crossref

33 Tesfaye Abera Gudeta, Tilahun Mekonnen Regassa, Alemayehu Sayih Belay. "Magnitude and factors associated with anemia among pregnant women attending antenatal care in Bench Maji, Keffa and Sheka zones of public hospitals, Southwest, Ethiopia, 2018: A cross-sectional study", PLOS ONE, 2019

11 words — < 1%

Crossref

34	docz.io Internet	11 words — < 1%
35	pt.scribd.com Internet	11 words — < 1%
36	www.researchgate.net Internet	11 words — < 1%
37	Catarina Bourgard, Letusa Albrecht, Ana C. A. V. Kayano, Per Sunnerhagen, Fabio T. M. Costa. "Plasmodium vivax Biology: Insights Provided by Genomics, Transcriptomics and Proteomics", <i>Frontiers in Cellular and Infection Microbiology</i> , 2018 Crossref	10 words — < 1%
38	M. Mutalazimah, Diah Pitaloka Putri. "Helminthiasis, iron intake, and hemoglobin levels in pregnant women", <i>Enfermería Clínica</i> , 2021 Crossref	10 words — < 1%
39	core.ac.uk Internet	10 words — < 1%
40	etheses.whiterose.ac.uk Internet	10 words — < 1%
41	onlinelibrary.wiley.com Internet	10 words — < 1%
42	proceeding.tenjic.org Internet	10 words — < 1%
43	www.ajtmh.org Internet	10 words — < 1%

44	www.ejurnalmalahayati.ac.id Internet	10 words — < 1%
45	www.researchsquare.com Internet	10 words — < 1%
46	Berhanu Senbeta Deriba, Elias Teferi Bala, Gizachew Abdissa Bulto, Tinsae Abeya Geleta et al. "Determinants of Anemia among Pregnant Women at Public Hospitals in West Shewa, Central Ethiopia: A Case-Control Study", Anemia, 2020 Crossref	9 words — < 1%
47	Octa Dwienda Ristica. "Faktor Risiko Kejadian Anemia pada Ibu Hamil", Jurnal Kesehatan Komunitas, 2013 Crossref	9 words — < 1%
48	Wiley, Andrea S.. "Medical Anthropology", Oxford University Press Publications	9 words — < 1%
49	Yirga Ebuy, Mussie Alemayehu, Mengistu Mitiku, Gelila K. Goba. "Determinants of severe anemia among laboring mothers in Mekelle city public hospitals, Tigray region, Ethiopia", PLOS ONE, 2017 Crossref	9 words — < 1%
50	annalsofrscb.ro Internet	9 words — < 1%
51	ecronicon.com Internet	9 words — < 1%
52	iiste.org Internet	9 words — < 1%

53	pdfs.semanticscholar.org Internet	9 words — < 1%
54	prp.hec.gov.pk Internet	9 words — < 1%
55	researchspace.ukzn.ac.za Internet	9 words — < 1%
56	worldwidescience.org Internet	9 words — < 1%
57	www.science.gov Internet	9 words — < 1%
58	Emily C. Keats, Christina Oh, Tamara Chau, Dina S. Khalifa, Aamer Imdad, Zulfiqar A. Bhutta. "Effects of vitamin and mineral supplementation during pregnancy on maternal, birth, child health and development outcomes in low - and middle - income countries: A systematic review", Campbell Systematic Reviews, 2021 Crossref	8 words — < 1%
59	bmchematol.biomedcentral.com Internet	8 words — < 1%
60	e-journal.unair.ac.id Internet	8 words — < 1%
61	eprints.keele.ac.uk Internet	8 words — < 1%
62	horizon.documentation.ird.fr Internet	8 words — < 1%
63	repository.um-surabaya.ac.id Internet	8 words — < 1%

8 words — < 1%

64 www.jpmmh.org
Internet

8 words — < 1%

65 Tesema Abera Yada, Yadeta Dessie, Rasha Darghawth, Tara Wilfong, Mohammed Abdurke Kure, Kedir Teji Roba. "Magnitude of Intestinal Parasitosis, Malnutrition, and Predictors of Anemia Among Nonpregnant Reproductive-Age Women Attending Healthcare Services in Olenchity General Hospital, Central Ethiopia", *Frontiers in Tropical Diseases*, 2021

7 words — < 1%

Crossref

66 journal.umpalangkaraya.ac.id
Internet

7 words — < 1%

67 Belayneh E Dejene, Tesfamariam M Abuhay, Dawit S Bogale. "Predicting the Level of Anemia among Ethiopian Pregnant Women using Homogeneous Ensemble Machine Learning Algorithm", *Research Square Platform LLC*, 2022

6 words — < 1%

Crossref Posted Content

68 Devaraj Acharya, Ramesh Adhikari, Padam Simkhada. "Prevalence and Determinants of Anaemia among Women aged 15-49 in Nepal: A Trend Analysis from Nepal Demographic and Health Surveys from 2006 to 2016", *Asian Journal of Population Sciences*, 2022

6 words — < 1%

Crossref

69 Li Lin, Yumei Wei, Weiwei Zhu, Chen Wang, Rina Su, Hui Feng, Huixia Yang. "Prevalence, risk factors and associated adverse pregnancy outcomes of anaemia in Chinese pregnant women: a multicentre retrospective study", *BMC Pregnancy and Childbirth*, 2018

6 words — < 1%

70 Lucas Gosdin, Reynaldo Martorell, Rosario M. Bartolini, Rukshan Mehta, Sridhar Srikantiah, Melissa F. Young. "The co - occurrence of anaemia and stunting in young children", Maternal & Child Nutrition, 2018

6 words — < 1%

Crossref

EXCLUDE QUOTES OFF

EXCLUDE SOURCES OFF

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF