

# OVERVIEW OF NURSING PROBLEM IN INPATIENTS WITH DRUG-RESISTANT TUBERCULOSIS AT SULIANTI SAROSO INFECTIOUS DISEASE HOSPITAL

*By Maemun et al*

**Original Research Article**

## NURSING PROBLEM IN INPATIENTS WITH DRUG-RESISTANT TUBERCULOSIS AT SULIANTI SAROSO INFECTIOUS DISEASE HOSPITAL

Veronika Hutabarat<sup>1</sup>, Farida Murtiani<sup>1</sup>, Nuraidah<sup>1</sup>, Yuni Shahroh<sup>1</sup>, Sumaryati<sup>1</sup>, Ramlah<sup>1</sup>, Hasinah<sup>1</sup>, Kunti Wijiarti<sup>1</sup>, Aninda Dinar Widiantari<sup>1</sup>, Siti Maemun<sup>1,2\*</sup>

<sup>1</sup>Rumah Sakit Penyakit Infeksi Prof. Dr. Sulianti Saroso  
<sup>2</sup>Fakultas Ilmu Kesehatan, Universitas Respati Indonesia

\*Correspondence:

**Siti Maemun**

Fakultas Ilmu Kesehatan, Universitas Respati Indonesia

Bambu Apus, Cipayang, Jakarta, 13890, Indonesia

Email: [maumun83@gmail.com](mailto:maumun83@gmail.com)

**Article Info:**

Received: April, 5<sup>th</sup>, 2024

Revised: June, 2<sup>nd</sup>, 2024

Accepted: August, 24<sup>th</sup>, 2024

**DOI:**

<https://doi.org/10.36720/nhjk.v14i1.647>

### Abstract

**Background:** Tuberculosis is a worldwide health problem, including in Indonesia. Drug-resistant tuberculosis (DR-TB) is caused by spontaneous mutations in the chromosomes that cause resistance to first-line DR-TBugs.

**Objectives:** to determine the patient characteristics and nursing problems of patients with DR-TB.

**Methods:** We conducted a cross-sectional study of inpatients with DR-TB in the DR-TB isolation ward at Sulianti Saroso Infectious Disease Hospital from July to December 2021. The variables investigated in this study include patient characteristics, comorbid diseases, tuberculosis treatment history, nutritional status, psychological status, nursing problem, and subcategories of nursing problems.

**Results:** Most patients were 56-65 years old, male, high school graduates, unemployed, and married. Most of the populations had diabetes mellitus as a comorbid disease, a history of having received previous treatment, and long-term treatment regimens. Most patients had normal nutritional status and a calm and adaptive psychological history. Nursing problems were airway clearance, dietary deficits, and electrolyte imbalance. The sub-categories of nursing problems that appear most were nutrition and fluid intake, respiration, and ego integrity. This occurs due to physiological disturbances caused by the disease process and the side effects of DR-TBugs.

**Conclusion:** The process of *Mycobacterium tuberculosis* infection accelerates the body's metabolism as it combats the bacteria. This resulted in decreased appetite and reduced nutritional status. Consequently, nurses should focus on intensified care for DR-TB patients. This involves ensuring adequate nutrition and fluid intake, offering spiritual support to boost confidence, and educating patients on maintaining airway hygiene.

**Keywords:** *Nursing, Resistance, Treatment*



## INTRODUCTION

Drug-resistant tuberculosis (DR-TB) is one of the major global health issues. DR-TB is caused by spontaneous mutation in chromosomes, leading to resistance against first-line anti-tuberculosis drugs (ANTI-TB DRUGS). In 2020, it was estimated that there were 10 million TB cases, with 0.5 million of them being MDR TB/DR-TB (WHO 2021). In Indonesia, out of 824,000 tuberculosis (TB) cases in 2020, there were 7,921 cases of drug-resistant tuberculosis (DR-TB), with 4,590 still undergoing treatment (Kemenkes, 2022).

Inadequate treatment can cause bacterial resistance to anti-tuberculosis drugs (anti-TB drugs). This can be caused by several factors, including incorrect diagnosis by healthcare workers, inappropriate dosages, types, quantities, and duration of treatment, and limited patient education (Kemenkes 2020). The role of nurses is crucial in reducing drug-resistant tuberculosis (DR-TB) cases.

According to Law Number 38 of 2014, nursing care is a series of interactions between nurses, clients, and their environment (Indonesia 2014). This aims to empower clients to care for themselves and meet their needs. Nursing care includes five consecutive stages: assessment, diagnosis, intervention, implementation, and evaluation. According to Zori & Morrison, a nurse's competency in identifying patient problems and determining appropriate interventions depends on the sharpness of analysis and critical thinking following the evidence base (Zori and Morrison, 2009).

Nursing problems are one of the main components in nursing diagnosis and describe the essence of the patient's response to their health condition. Nursing problems are established based on the grouping of data and interpretation of data obtained from the patient's nursing assessment. Nursing assessment includes subjective data (patient complaints) and objective data (inspection, percussion, palpation, and auscultation) (Anggi 2018; Leniwita and Anggraini 2019).

The next step after nursing assessment is to identify problems and determine nursing issues such as patterns of nutritional metabolism, exercise activity patterns, and health management perceptions. Determining nursing problems in TB-DR patients is crucial for planning appropriate actions to address these nursing issues, thus preventing medication non-adherence and discontinuation of anti-TB drug treatment.

Often, nurses are the first point of contact when patients seek treatment and serve as cadres in their role as healthcare providers who administer or supervise the daily medication of patients. Nurses are also frequently the first to hear complaints about drug side effects from patients and, therefore, can promptly provide interventions. Thus, efforts are needed to help nurses identify symptoms that may indicate tuberculosis drug side effects; assess the severity of drug side effects and the possibility of other causes; and act appropriately to alleviate patient discomfort, reduce the progression of drug side effects, and support patients in completing their treatment (Raftery et al. 2018).

Based on a case study conducted at the referral hospital, there are five nursing problems in DR-TB patients: impaired gas exchange, ineffective airway clearance, imbalance nutrition, deficit knowledge, and sudden/acute pain related to biological sensory agents (Neni Lestari 2020; Sri Okta Novelia 2019). In addition, according to Rifqi Arafat's research (2019), nursing problems found in DR-TB patients include ineffective breathing patterns, ineffective health management (limited therapeutic knowledge), and ineffective family health management (complexity of therapeutic regimen) (Arafat, 2019).

Sulianti Suroso Infectious Diseases Hospital (SSIDH) is a referral centre for drug-resistant tuberculosis (DR-TB) patients. In 2019, 120 DR-TB cases were treated. SSIDH has been designated as a referral hospital for the management of drug-resistant tuberculosis. The

high number of cases has led to various nursing problems that may differ between individual DR-TB patients. Therefore, nurses must identify common nursing problems to determine appropriate interventions for DR-TB patients and the relevance between literature and the implementation of nursing care in the field.

**Objective(s):** to determine the characteristics of DR-TB patients treated and to identify common nursing problems among DR-TB patients at SSIDH.

**METHOD**

*Study Design*

This research is a descriptive quantitative study with a cross-sectional design.

*Setting*

This research was conducted in July to December 2021 at Sulianti Saroso Infectious Disease Hospital.

*Research Subject*

The population of this study comprises all DR-TB patients admitted to Dahlia 1 Ward from July to December 2021 are 36 patients. We use total population in this study. The sample consists of all inpatients diagnosed with DR-TB, >18 years old, and hospitalized once during the period from July to December 2021. The data source is secondary data from medical records, including patient characteristics, comorbidities, nutritional status, treatment history, subjective/objective data assessment, and nursing problems according to categories and sub-categories.

*Instruments*

The data collection instrument uses a Case Report Form (CRF). The measurement of the variables, nursing problems and subcategories of nursing problems, is based on the findings of the nursing assessment in the form of subjective and objective data. This data then analyzed to determine the nursing problems and subcategories of nursing

problems. The classification of nursing problems and subcategories of nursing problems is based on the Indonesian Nursing Diagnosis Standard.

Based on subjective and objective data, they are categorized into major and minor. If 80%-100% major categories are found, then the nursing problem is actual. If less than 80%, then the nursing problem is a risk.

*Data Analysis*

Descriptive analysis will be conducted on all research variables in frequency (n) and percentage (%).

*Ethical Consideration*

This research has obtained ethical clearance from the Research Ethics Committee of SSIDH with approval number 40/XXXVIII.10/VII/2022.

**RESULT**

The research findings indicate that the common age group among respondents is 56-65, with the youngest being 19 and the oldest being 71. Most patients are male, have completed high school education, are unemployed, married, and reside in the Jakarta Bogor Depok Tangerang Bekasi (Jakarta Metropolitan Area) (Table 1).

**Table 1. Patient Characteristics (n=36)**

Sociodemographic	n	%
<b>Age (Year)</b>		
>18-25	6	16.7
26-35	5	13.9
36-45	7	19.4
46-55	7	19.4
56-65	10	27.8
66-75	1	2.8
<b>Gender</b>		
Male	21	58.3
Female	15	41.7
<b>Level of Education</b>		
Incomplete Primary School	3	8.3

Elementary School or Equivalent	6	16.7
Junior High School or Equivalent	7	19.4
High School or Equivalent	19	52.8
Diploma/Bachelor's Degree	1	2.8
<b>Employment Status</b>		
Unemployed	14	38.9
Daily labourer	4	11.1
Employee	12	33.3
Governance/Military Army/Police	5	13.9
Entrepreneur	1	2.8
<b>Marital Status</b>		
Married	26	72.2
Single	8	22.2
Divorced	2	5.6
<b>Domicile</b>		
Jakarta Metropolitan Area (JMA)	33	91.7
Outside JMA	3	8.3

Table 2 shows the medical history of TB-DR patients. The majority have comorbidities, the most common being diabetes mellitus (DM). Regarding treatment history, most have received previous treatment, with most receiving long-term treatment regimens. In terms of nutritional status, the majority are of normal weight. Based on psychological history, most are calm and adaptive.

**Table 2. Comorbid Disease History (n=36)**

Comorbid Diseases	N	%
<b>Presence of Comorbidities</b>		
Yes	21	58.3
No	15	41.7
<b>Type of Comorbidities</b>		
DM	15	41.7
Hypertension	3	8.3
HIV	14	38.9
Hepatitis B	2	5.6
Heart disease	1	2.8
Autoimun	1	2.8

<b>History of Previous Treatment</b>		
Yes	22	61.1
Discontinued	10	27.8
Never	4	11.1
<b>Regimen</b>		
Long-Term	35	97.2
Short-Term	1	2.8
<b>Nutritional Status</b>		
Undernourished	15	41.7
Normal	20	55.6
Overnourished	1	2.7
<b>Psychological</b>		
Anxious	16	44.4
Calm and Adaptive	18	50.0
Not Assessed	2	5.6

The most common nursing issues are airway clearance, nutritional deficit, and electrolyte imbalance. All of patients have more than one nursing problems (Table 3).

**Table 3. Nursing Problems**

Nursing Problems	N	%
Airway Clearance	23	26.1
Impaired Gas Exchange	1	1.1
Ineffective Breathing Pattern	5	5.7
Nutritional Deficit	20	22.7
Electrolyte Imbalance	10	11.4
Blood Glucose Instability	8	9.1
Activity Intolerance	1	1.1
Impaired Peripheral Perfusion	5	5.7
Anxiety	9	11.4
Risk of Infection	1	1.1
Discomfort/Pain Disorder	2	2.3
Acute Pain	1	1.1
Skin Integrity Impairment	1	1.1

The most frequent subcategories of nursing problems are nutrition and fluid, respiratory, and ego integrity. The majority of patients have more than subcategories of nursing problems (Table 4).

**Table 4. Subcategories of Nursing Problems**

Subcategories of Nursing Problems	N	%
Respiration	23	29.9
Circulation	6	7.8
Nutrition and Fluid	31	40.3
Activity and Rest	1	1.3
Safety and Protection	2	2.6
Pain and Comfort	3	3.9
Ego Integrity	11	14.3

## DISCUSSION

### Patient Characteristics

More patients are at 56-65 years old by range (19-71 years old). Meanwhile, a study by Manggasa and Suharto (2022) showed that DR-TB patients ranged from 16 to 30 years old (Manggasa and Suharto, 2022). Another study states that most patients fall within the age range of 18-40 years (80%), while those above 40 years old account for 20% (Wibowo, Burhan, and Putra 2021). The study by Munawwarah et al., cited in Tri Wahyuni, mentions that age significantly influences the higher resistance to anti-TB drugs, with the highest resistance observed among the age group of 31-40 years, at 46.7% (Wahyuni 2020).

The highest incidence of MDR-TB patients occurs in the productive age group because, during this period, there is frequent social interaction with others and high mobility, consequently increasing the risk of transmission to others and the surrounding environment (Azwar, Noviana, and Hendriyoto 2017). Furthermore, non-compliance with medication increases the risk of resistance (Horter et al. 2014). The WHO report states that most tuberculosis patients in

developing countries are in the working age group (15-54 years old).

In this study, the majority of DR-TB cases are found in males. Similar findings are reported in Dr. H. Chasan Bosoirie Terrate Hospital, where male patients outnumber female patients (Bayan, Prihanto, Anwar 2022; Manggasa and Suharto 2022). The WHO report indicates that the prevalence of TB in males is 1.7 times higher than in females. This is attributed to biological, cultural, and social factors, where males often engage in activities outside the home more frequently than females.

The risk of the disease is not influenced by gender, but males tend to be at higher risk due to smoking behaviour. Patients with a history of smoking are more susceptible to TB infection compared to non-smokers (Anisah, Sumekar, and Budiarti 2021). Furthermore, other research (Chuchottaworn et al. 2015) suggests that adherence to anti-TB drugs among male patients is relatively not lower than that of females, hence the higher prevalence of MDR-TB among males (Chuchottaworn et al. 2015).

Based on education, the majority of DR-TB patients have completed secondary education, specifically high school. In a study conducted at the Umbulharjo Primary Health Center in Yogyakarta, most respondents, family members of pulmonary TB patients, graduated from high school or equivalent (50.0%). Secondary education makes it easier to receive information, thus influencing thoughts, actions, and increasing knowledge (Notanti-TB drugsmojo 2018).

This research proves that the majority of DR-TB patients have higher education levels, but their knowledge about the risk factors for DR-TB is minimal. Additionally, there is a lack of awareness regarding medication adherence and completing the treatment regimen (Bijawati, Amansyah, and Nurbiah, 2018). Having a higher education level does not guarantee that someone's knowledge about a particular topic, including the occurrence of DR-TB, will be better than those with lower

education. However, exposure to information also plays a significant role in shaping one's knowledge.

This study's results indicate that most DR-TB patients are unemployed. This is consistent with the findings at Dr. H. Chasan Boesoerie Ternate Hospital, where most DR-TB patients are also unemployed (31.8%) (Bayan et al. 2022). Flora et al.'s study on the risk factors of DR-TB in Bangladesh reported that 51 controls (33.6%) and 46 cases (34.1%) are unemployed (Flora et al. 2013).

Occupation is closely related to income, which impacts purchasing power in meeting daily needs, including nutritional requirements. Adequate nutrition strongly influences nutritional status through parameters such as BMI. Nutritional status affects the body's immunity, which makes individuals vulnerable to diseases, including tuberculosis (A Dwi Sarwani; Nurfaela, Sri 2012).

#### Medical History

Most DR-TB patients have comorbid DM, and this finding is consistent with research in China, which shows that DR-TB patients with DM are more prevalent, at 17.7%, compared to DR-TB patients without DM, which is 9.3% ( $p$ -value  $<0.01$ ). According to Zhang Qing et al., DM can worsen TB infection, prolong treatment duration, reduce medication adherence, and increase the frequency of treatment side effects. Irregular DM medication intake leads to uncontrolled blood sugar levels, making TB infection challenging to cure. Also, due to low immunity, DM patients may become re-infected with DR-TB strains. TB-DM cases may also have a poor prognosis if relapse occurs within two years, requiring new DM therapy and effective treatment for DR-TB patients with DM (Fauziah 2013).

A systematic review states that patients with DM are at a higher risk of experiencing MDR compared to those without DM because DM causes lower immunity, leading to a more severe prognosis of tuberculosis infection

(Restinia et al., 2021). Additionally, the results are more lung inflammation (Utomo, Nugroho, and Margawati 2016).

Based on treatment history, the majority have received previous treatment. This finding is consistent with the research by Triandari & Rahayu (2018), which suggests a relationship between the history of prior TB treatment outcomes and DR-TB. Patients with a history of treatment discontinuation are three times more likely to experience DR-TB compared to patients without a history of treatment discontinuation. Therefore, the treatment history is the most dominant factor in the occurrence of drug resistance. Treatment failure and discontinuation are causes of DR-TB (Triandari and Rahayu 2018) and MDR-TB (Manggasa and Suharto 2022). Having more than one history of TB treatment increases the risk of DR-TB occurrence, which is related to previous treatment outcomes, such as treatment discontinuation, treatment failure, relapse, or patients who have undergone DR-TB treatment.

DR-TB means the TB bacteria are resistant to drugs that can kill them, so the principle of treating DR-TB is to use antibiotics that are still expected to kill the TB bacteria within 9-24 months, depending on which drugs the patient can use. Treatment is divided into two types: Short Treatment Regimen (STR), which requires a minimum of 9 months, and Individual Treatment, which involves a minimum of 20 months. Each treatment is divided into two stages: the initial phase, which lasts about 4-6 months, and the continuation phase (Dinas Kesehatan Provinsi NTB 2023).

The results indicate that the majority are receiving long-term treatment regimens. Treating DR-TB requires a longer duration (18-24 months) compared to regular TB (6-8 months). This is because the bacteria causing TB are already resistant to first-line anti-TB drugs, so treating DR-TB requires more potent anti-TB drugs and often involves more injections and medication (Aviana, Juti, and Budiyantri 2021).

The previous treatment regimen for DR-TB required a long period of treatment. Therefore, in May 2016, WHO recommended a new DR-TB treatment regimen involving a short-term guideline. This new regimen aims to make the treatment period more effective, thus reducing the overall treatment duration and minimising treatment interruption. With a prolonged treatment period, DR-TB patients are at risk of treatment interruption, which can ultimately lead to treatment failure. This recommendation is based on observational studies conducted in countries that have implemented short-term treatment regimens, including Bangladesh, Benin, Burkina Faso, Burundi, Cameroon, Central Africa, Congo, Niger, Swaziland, and Uzbekistan. WHO reports show that short-term treatment success rate is higher than long-term regimens. Shorter treatment durations and more effective treatment outcomes are expected to increase treatment enrollment, reduce interruption rates, and improve success rates for DR-TB patients in Indonesia (Agustina, Maulida, and Yovsyah 2018).

Based on their nutritional status, most subjects are in the normal weight category. Other research indicates that the majority of their study populations are still within the normal range (49.2%) (Wibowo et al. 2021) (Santy 2020). In contrast, a study in India found that the majority of TB patients had low body mass index (BMI) (Bhargava et al. 2013), which can be a risk factor for drug resistance. Nutritional status plays a crucial role in tuberculosis treatment, as good dietary status leads to better treatment response. BMI is used as an indicator to assess nutritional status. Increasing body weight during treatment can accelerate the improvement of nutritional status, leading to faster sputum conversion and clinical improvement of the lung (Wibowo et al. 2021).

Based on their psychological history, most patients are calm and adaptive (50%) and anxious (44.4%). Acceptance of the disease becomes a problem for DR-TB patients. This

issue is prevalent among adults and older people and affects teenagers. Many of them struggle to adjust to their condition and face limitations in carrying out daily activities, which often leads to feelings of despair. Therefore, self-acceptance and support from those around them can help patients cope with their daily lives.

#### Nursing Problems

The most prevalent nursing problem in DR-TB cases is ineffective airway clearance. This finding is consistent with research by Rifki (2019) and SO Novela (2019), which also identified ineffective airway clearance as the primary issue in DR-TB patients. This is due to the mycobacterium tuberculosis bacteria attacking the lungs through the inflammatory process in the alveoli, leading to excessive sputum buildup and resulting in problems with ineffective airway clearance (Nararif and Kusuma 2015). According to SDKI in 2016, airway clearance problems are caused by airway spasms, hypersecretion of the airway, and retained secretions characterised by excessive coughing with sputum. Airway clearance is the second-highest nursing problem after nutrition and fluid-related concerns.

A prospective study conducted at Helmi Chest Hospital found that, in comparison to routine nursing care, targeted nursing interventions combined with psychological counseling significantly improve the quality of life and reduce negative emotions in patients with XDR-TB, as well as decrease the incidence of complications (Lu et al. 2021).

Nutritional deficit is also a significant nursing problem in TB MDR cases, accounting for 22.73%. According to Garba Garuda's 2020 analysis, the most prevalent dietary status among TB patients, as measured by Body Mass Index (BMI), is underweight (50.27%). This is due to the *Mycobacterium tuberculosis* infection process, which increases metabolism in the body as it fights against the bacteria. Consequently, this increased metabolism leads

to side effects such as decreased appetite and reduced nutritional status. Nutritional deficit nursing problems are caused by increased metabolic needs and the inability to digest food and are defined as a weight loss of more than ten per cent of the ideal range. The most prevalent subcategory of nursing problems is nutrition and fluid-related concerns (40.26%). This is attributed to physiological disturbances due to the disease process and the side effects of DR-TB medications.

Based on the research findings, the nursing problem of the risk of electrolyte imbalance is the highest. Reviono's study (2014) demonstrated that DR-TB patients experience hypokalemia (52.6%), nausea (79.8%), and vomiting (78.9%) (Reviono et al. 2014). Syefi Nuraeni Fitriana (2023) stated that 70.45% of DR-TB patients receiving kanamycin treatment experienced hypokalemia (Fitriana, Zulkarnain, and Yulimanida, 2023). The causes of electrolyte imbalance are multifactorial, possibly due to drug side effects and gastrointestinal disturbances. The subcategory of nursing problems related to psychological issues is ego integrity, which is caused by high levels of anxiety, helplessness, and despair. The high prevalence of ego integrity problems is attributed to the side effects of medication.

These findings provide insights for nurses to provide more intensive care for DR-TB patients, especially regarding nutritional and fluid management, religious care to foster belief in recovery, and educating patients on maintaining airway clearance independently.

## CONCLUSION

Based on the medical history of DR-TB patients, the majority have the highest comorbidity of DM, have received or previously received treatment, are undergoing long-term treatment regimens, have normal weight nutritional status, and have calm and adaptive psychological histories. The most prevalent nursing problems are airway clearance, nutritional deficit, and electrolyte

imbalance. The most pervasive subcategories of nursing problems are nutrition and fluid-related concerns, respiration, and ego integrity.

## RESEARCH LIMITATIONS

This study focuses on one referral hospital for DR-TB cases; therefore, its findings cannot be generalised. Further limitations include a small sample size. One reason for the small sample size is the COVID-19 pandemic in 2021, during which SSIDH only treated COVID-19 patients. In October-December 2021, inpatient services for DR-TB patients resumed.

## ACKNOWLEDGEMENT

We thank the Chief Executive Officer and Pulmonologists for granting research permission and the research unit for supporting the implementation of this study.

## FUNDING

There was no funding for this research.

## AUTHOR CONTRIBUTION

**Veronika Hutabarat**: Compiling research ideas, as a leader in the research process.

**Murtiani**: Assist in data processing.

**Nuraidah**: Assist in data processing.

**Yuni Shahroh**: Assist in data processing.

**Sumaryati**: Assist in data processing.

**Ramlah**: Assist in data processing.

**Hasinah**: Assist in data processing.

**Kunfi Wijarti**: Assist in data processing.

**Aninda Dinar Widiantari**: Assist in data processing.

**Siti Maemun**: Assist in the formation of research ideas.

**ORCID**

**Veronika Hutabarat:**

<https://orcid.org/0009-0004-9473-7819>

28

**Murtiani:**

<https://orcid.org/0000-0003-4992-117X>

**Armaidah:**

<https://orcid.org/0000-0001-8083-5443>

**Yuni Shahroh:**

<https://orcid.org/0009-0003-1379-8234>

**Sumaryati:**

<https://orcid.org/0009-0002-3920-5540>

**Ramlah:**

<https://orcid.org/0009-0004-4108-6311>

**Hasinah:**

<https://orcid.org/0009-0002-1407-9971>

**Kunti Wijarti:**

<https://orcid.org/0000-0002-8932-5041>

**Inda Dinar Widiantari:**

<https://orcid.org/0000-0002-5376-6208>

**Siti Maemun:**

<https://orcid.org/0000-0001-8841-9591>

**DAFTAR PUSTAKA**

A Dwi Sarwani, Nurfaela, Sri, Isnani Zahrotul, S. R. 2012. "Faktor Risiko Multidrug RESISTANT TUBERCULOSIS (MDR-TB)." *Jurnal Kesehatan Masyarakat* 8(Vol 8, No 1 (2012)):60-66.

Agustina, Rina, Rizka Maulida, and Yovsyah. 2018. "Faktor-Faktor Yang Berhubungan Dengan Kesuksesan Kesembuhan Dari Pengobatan Regimen Pendek ( Short Treatment Regimen ) Pada Pasien Tuberkulosis Resistensi Obat Di Indonesia Tahun 2017 Factors Associated with Recovery Success after Short Treatment Regimen." *Jurnal*

*Epidemiologi Kesehatan Indonesia* 2(2):65-71.

Anggi, Pebriana. 2018. "Pengkajian Dalam Proses Keperawatan Anamnesa Dan Pemeriksaan Fisik Abstrak Latar Belakang." *Pengkajian Dalam Proses Keperawatan Anamnesa Dan Pemeriksaan Fisik* 11.

Anisah, Anisah, Dyah Wulan Sumekar, and Endang Budiarti. 2021. "Hubungan Demografi Dan Komorbid Dengan Kejadian Tuberkulosis Resistensi Obat (TB RO)." *Jurnal Ilmiah Kesehatan Santri Husada* 10(2):568-74. doi: 10.35816/jiskh.v10i2.655.

Arafat, Rifqi. 2019. "Asuhan Keperawatan Tn. D.U Dengan MDR-TB (Multi Drug Resistance-Tuberculosis Paru) Di Ruang Tulip RSUD Prof. Dr. W. Z. Johannes Kupang." *Jurnal Keperawatan*.

Aviana, Fitri, Sutopo Patria Jati, and Rani Tiya Budiyanti. 2021. "Systematic Review Pelaksanaan Programmatic Management of Drug-Resistant Tuberculosis Pada Pasien Tuberkulosis Resistensi Obat." *Jurnal Kesehatan Masyarakat (Undip)* 9(2):215-22. doi: 10.14710/jkm.v9i2.28719.

Azwar, Gusti Andhika, Dewi Indah Noviana, and FX. Hendriyono. 2017. "Karakteristik Penderita Tuberkulosis Paru Dengan Multidrug-Resistant Tuberculosis (Mdr-Tb) Di Rsud Ulin Banjarmasin." *Berkala Kedokteran* 13(1):23. doi: 10.20527/jbk.v13i1.3436.

Bayan, Uways Al-Qarni, Eko Sudarmo D, Prihanto, and Mawardi Anwar. 2022. "Karakteristik Pasien Tuberkulosis Resistensi Obat Di RSUD Dr. H. Chasan Boesoirie." *Kieraha Medical Journal* 4(1):15-21.

Bhargava, Anurag, Madhuri Chatterjee, Yogesh Jain, Biswaroop Chatterjee, Anju Kataria, Madhavi Bhargava, Raman Kataria, Ravi D'Souza, Rachna Jain, Andrea Benedetti, Madhukar Pai, and Dick Merzies. 2013. "Nutritional Status

- of Adult Patients with Pulmonary Tuberculosis in Rural Central India and Its Association with Mortality." *PloS One* 8(10):e77979. doi: 10.1371/journal.pone.0077979.
- Bijawati, Emni, Munawir Amansyah, and Nurbiah. 2018. "The Risk Factors for Treatment of Multidrug Resistance Tuberculosis (MDR-TB) Patients in Labuang Baji General Hospital Makassar in 2017." *Jurnal Nasional Ilmu Kesehatan* 1:1–17.
- Chuchotaworn, Charoen, Vipa Thanachartwet, Piamlarp Songsayanh, Thu Zar Myint Thun, Duangjai Sahassananda, Manoon Surabotsophon, and Varanee Desakorn. 2015. "Risk Factors for Multidrug-Resistant Tuberculosis among Patients with Pulmonary Tuberculosis at the Central Chest Institute of Thailand." *PloS One* 10(10):e0139986. doi: 10.1371/journal.pone.0139986.
- Dinas Kesehatan Provinsi NTB. 2023. "Tuberkulosis Resisten Obat, Bukan Tuberkulosis Biasa." *Datin Litbangkes*.
- Fauziah, L.A. 2013. "Faktor-Faktor Yang Berpengaruh Terhadap Kejadian Tuberculosis Multidrug Resisten(TB-MDR) Di RSUP Persahabatan." *Departemen Epidemiologi FKM UI*.
- Fitriana, Syefi Nuraeni, Bambang Subukti Zulkarnain, and Herri Yulimankita. 2023. "Analisis Efek Samping Kanamisin Dan Kapreomisin Terhadap Fungsi Ginjal Pada Pasien TB RO Di RSUD Dr. Soetomo." *Cendekia Journal of Pharmacy* 7(1): 10–19.
- Flora, M. S., M. N. Amin, M. R. Karim, S. Afroz, S. Islam, A. Alam, and M. Hossain. 2013. "Risk Factors of Multi-Drug-Resistant Tuberculosis in Bangladeshi Population: A Case Control Study." *Bangladesh Medical Research Council Bulletin* 39(1):34–41. doi: 10.3329/bmrcb.v39i1.15808.
- Horter, Shona, Beverley Stringer, Lucy Reynolds, Muhammad Shoaib, Samuel Kasozi, Esther C. Casas, Meggy Verputten, and Philipp da Cros. 2014. "'Home Is Where the Patient Is': A Qualitative Analysis of a Patient-Centred Model of Care for Multi-Drug Resistant Tuberculosis." *BMC Health Services Research* 14:81. doi: 10.1186/1472-6963-14-81.
- Indonesia, Pemerintah Pusat. 2014. *Undang-Undang (UU) Nomor 38 Tahun 2014 Tentang Keperawatan*. Indonesia.
- Kemendes. 2022. *Laporan Tahunan 2021*. Jakarta.
- Kemendes, P2PL. 2020. *Petunjuk Teknis Penatalaksanaan Tuberculosis Resistan Obat Di Indonesia*.
- Leniwita, Husain, and Yanti Anggraini. 2019. "Modul Dokumentasi Keperawatan." *Universitas Kristen Indonesia* 1–182.
- Lu, Yanhua, Hongqian Wang, Jianfeng Zhu, Na Wang, Dan Cai, and Lingdi Li. 2021. "Effect of Targeted Nursing Intervention plus Psychological Counseling on Quality of Life, Negative Emotions, and Complications in Patients with Extensively Drug-Resistant Tuberculosis." *American Journal of Translational Research* 13(12):13950–58.
- Manggasa, Dafrosia Darmi, and Dewi Nurviana Suharto. 2022. "Riwayat Pengobatan Dan Komorbid Diabetes Mellitus Berhubungan Dengan Kejadian Tuberculosis Resisten Obat." *Poltekita: Jurnal Ilmu Kesehatan* 15(4):403–8. doi: 10.33860/jik.v15i4.659.
- Neni Lestari. 2020. "Asuhan Keperawatan Pada Pasien Dengan TB MDR Di Ruang Rawat Inap Pejuang Rumah Sakit Daerah Bangkinang." *Poltekkes Kemenkes Riau*.
- Notoatmojo. 2018. *Pendidikan Dan Perilaku Kesehatan*. Jakarta: PT Rineka Cipta.
- Nurarif, Amin Huda, and Hardhi Kusuma. 2015. *Asuhan Keperawatan Berdasarkan Diagnosa Medis & NANDA NIC NOC Jilid 3*. Edisi 3. Jogjakarta: Mediacion publishing.

- Raftery, Ann, Carrie Tudor, Lisa True, and Cattalina Navarro. 2018. *Panduan Keperawatan Untuk Manajemen Efek Samping Pengobatan Tuberkulosis Resistan Obat.*
- Restinia, Mita, Sondang Khairani, and Reise Manninda. 2021. "Faktor Resiko Penyebab Multidrug Resistant Tuberculosis: Sistematis Review." *Pharmaceutical and Biomedical Sciences Journal (PBSJ)* 3(1):9-16. doi: 10.15408/pbsj.v3i1.20049.
- Reviono, P. Kusnanto, Vicky Eko, Helena Pakiding, and Dyah Nurwidiasih. 2014. "Multidrug Resistant Tuberculosis (MDR-TB): Tinjauan Epidemiologi Dan Faktor Risiko Efek Samping Obat Anti Tuberculosis." *Majalah Kedokteran Bandung* 46(4):189-96. doi: 10.15395/mkb.v46n4.336.
- Santy, Diana. 2020. "Hubungan Indeks Massa Tubuh (Body Mass Index) Dengan Konversi Sputum Pada Pasien Tb Resisten Obat Yang Menjalani Pengobatan Dengan Panduan Jangka Pendek (Short Term Regimen) Di RSUP H. Adam Malik Medan, Sumatera Utara, Indonesia." Universitas Sumatra Utara.
- Sei Okta Novelia. 2019. "Asuhan Keperawatan Pada Pasien Dengan Multi Drug Resistance (TB MDR) Di Ruang Rawat Inap Pura RSUP Dr. M. Djamil Padang." Poltekkes Kemenkes RI Padang.
- Triandari, Devi, and Sri Rama Rahayu. 2018. "Kejadian Tuberculosis Multi Drug Resistant Di RSUP Dr. Kariadi." *Higea Journal of Public Health* 2(2):194-204.
- Utomo, Radityo, HS K. Heri Nugroho, and Ani Margawati. 2016. "Hubungan Antara Status Diabetes Melitus Tipe 2 Dengan Status Tuberculosis Paru Lesi Luas." *Jurnal Kedokteran Diponegoro* 5(4):1536.
- Wahyuni, Tri. 2020. "Faktor-Faktor Yang Berhubungan Dengan Kejadian Multidrug Resistant Tuberculosis (Mdr-Tb)." WHO. 2021. "Global Tuberculosis Report 2021." Retrieved January 25, 2024 (<https://www.who.int/publications/i/item/9789240037021>).
- Wibowo, Adityo, Erlina Buthan, and Andika Chandra Putra. 2021. "Pola Resistansi Kuman Tuberculosis Dan Regimen Pengobatan Pada Pasien Tuberculosis Resisten Obat Di Rumah Sakit Pusat Rujukan Respirasi Nasional Persahabatan Jakarta." *Jurnal Kedokteran Universitas Lampung* 5(1):1. doi: 10.23960/jkunila511-6.
- Zori, Susan, and Barbara Morrison. 2009. "Critical Thinking in Nurse Managers." *Nursing Economics* 27(2):75-79,98.

Cite this article as: Hutabarat, V., et al. (2025). Overview Of Nursing Problem in Inpatients with Drug-Resistant Tuberculosis at Suliarti Saroso Infectious Disease Hospital. *Nurse and Health: Jurnal Keperawatan*, 14 (1), 1-11. <https://doi.org/10.36720/nhjk.v14i1.647>

# OVERVIEW OF NURSING PROBLEM IN INPATIENTS WITH DRUG-RESISTANT TUBERCULOSIS AT SULIANTI SAROSO INFECTIOUS DISEASE HOSPITAL

ORIGINALITY REPORT

# 15%

SIMILARITY INDEX

## PRIMARY SOURCES

1	<a href="http://ejournal-kertacendekia.id">ejournal-kertacendekia.id</a> Internet	263 words — 6%
2	Sisilia Indriasari Widianingtyas, Etik Lusiani, Agung Kurniawan Saputra. "THE INFLUENCE OF CHILD GROWTH STATUS AND FAMILY RELATIONSHIPS ON THE SOCIAL-EMOTIONAL DEVELOPMENT OF PRESCHOOL-AGE CHILDREN", Nurse and Health: Jurnal Keperawatan, 2024 Crossref	29 words — 1%
3	<a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Internet	27 words — 1%
4	<a href="http://journals.plos.org">journals.plos.org</a> Internet	26 words — 1%
5	Lono Wijayanti, Nur Ainiyah. "THE EFFECT OF THE SKIN PERSONAL HYGIENE MODULES AS HEALTH EDUCATION MEDIA AGAINST KNOWLEDGE IN PREVENTION OF SKABIES", Nurse and Health: Jurnal Keperawatan, 2019 Crossref	24 words — 1%
6	<a href="http://scifes.fkm.ui.ac.id">scifes.fkm.ui.ac.id</a> Internet	22 words — 1%
7	<a href="http://pubmed.ncbi.nlm.nih.gov">pubmed.ncbi.nlm.nih.gov</a> Internet	18 words — < 1%

- 
- 8 Yu Deng, Luxue Xie, Li Wang, Yaokai Chen. "Psychoanalysis of COVID-19 Patient Narratives: A Descriptive Study", *Medicina*, 2023  
Crossref 15 words — < 1%
- 
- 9 Yunita Arliny, Muhammad Fadjat Muarif, Wilda Mahdani, Dewi Behtri Yanifitri. "Anti-TB Drug Side-Effects on the Treatment of Drug-Resistant Tuberculosis (DR-TB) in dr. Zainoel Abidin Hospital Banda Aceh", *Jurnal Respirologi Indonesia*, 2025  
Crossref 15 words — < 1%
- 
- 10 [journal.unnes.ac.id](http://journal.unnes.ac.id)  
Internet 14 words — < 1%
- 
- 11 Salma Hameed, Sajid Rashid Ahmad, Mohammad Aqeel ur Rahman, Humera Nazir, Irfan Ullah. "Drug resistance profile of Mycobacterium tuberculosis and predictors associated with the development of drug resistance", *Journal of Global Antimicrobial Resistance*, 2019  
Crossref 12 words — < 1%
- 
- 12 Annisa Diyanabila Indrasari, Prima Belia Fathana. "STUDI LITERATUR : PATOGENESIS DAN DIAGNOSIS TUBERKULOSIS RESISTEN OBAT", *Jurnal Medika Malahayati*, 2024  
Crossref 11 words — < 1%
- 
- 13 Sita Paudel. "Risk Factors of Multidrug-Resistant Tuberculosis", *International Journal of Applied Sciences and Biotechnology*, 2017  
Crossref 10 words — < 1%
- 
- 14 [www.frontiersin.org](http://www.frontiersin.org)  
Internet 10 words — < 1%
- 
- 15 Bocar Baya, Chad J. Achenbach, Bourahima Kone, Yacouba Toloba et al. "Clinical Risk Factors 9 words — < 1%

Associated with Multidrug-Resistant Tuberculosis (MDR-TB) in  
Mali", International Journal of Infectious Diseases, 2019

Crossref

16 Riska Rohmawati, Ratna Yunita Sari, Imamatul  
Faizah, Siti Nur Hasina. "COMBINATION BENSON  
RELAXATION AND MURROTAL ALQURAN TO DECREASE  
ANXIETY, STRESS, AND DEPRESSION HEMODIALYSIS PATIENTS:  
AN EXPERIMENTAL STUDY", Nurse and Health: Jurnal  
Keperawatan, 2024

Crossref

17 [ijicc.net](http://ijicc.net) 9 words — < 1%

Internet

18 [inajemd.pbperkeni.or.id](http://inajemd.pbperkeni.or.id) 9 words — < 1%

Internet

19 [www.researchgate.net](http://www.researchgate.net) 9 words — < 1%

Internet

20 [www.researchsquare.com](http://www.researchsquare.com) 9 words — < 1%

Internet

21 Martinson, Melissa L., and Nancy E. Reichman.  
"Socioeconomic Inequalities in Low Birth Weight in  
the United States, the United Kingdom, Canada, and Australia",  
American Journal of Public Health, 2016.

Crossref

22 Nahla Ahmad Rahami, Rosita Dewi, Angga Mardro  
Raharjo, Cholis Abrori, Yuli Hermansyah,  
Muhammad Ali Shodikin. "Diabetes Mellitus Associated with  
Sputum Conversion Time in Drug-Resistant Pulmonary TB  
Patients at Dr. Soebandi Regional General Hospital, Jember",  
Jurnal Respirasi, 2024

Crossref

- 
- 23 Soedarsono Soedarsono, Tutik Kusmiati, Ariani Permatasari. "Effect of Diabetes Mellitus on Renal and Audiology Toxicities in Patients with Drug-Resistant Pulmonary Tuberculosis", Archives of Clinical Infectious Diseases, 2021  
Crossref 8 words — < 1%
- 
- 24 dspace.univ-sba.dz  
Internet 8 words — < 1%
- 
- 25 ejournal.poltekkes-smg.ac.id  
Internet 8 words — < 1%
- 
- 26 es.scribd.com  
Internet 8 words — < 1%
- 
- 27 ir.haramaya.edu.et  
Internet 8 words — < 1%
- 
- 28 journal.uwks.ac.id  
Internet 8 words — < 1%
- 
- 29 repository.lppm.unila.ac.id  
Internet 8 words — < 1%
- 
- 30 repository.stikim.ac.id  
Internet 8 words — < 1%
- 
- 31 sciendo.com  
Internet 8 words — < 1%
- 
- 32 Lailatul Fitriya, Kurnia Dwi Artanti. "TREATMENT OUTCOMES OF MULTIDRUG RESISTANT TUBERCULOSIS PATIENTS IN EAST JAVA FROM 2014 TO 2017", Jurnal Berkala Epidemiologi, 2020  
Crossref 7 words — < 1%

---

33 Irfan Ullah, Arshad Javaid, Zarfishan Tahir, Obaid Ullah, Aamer Ali Shah, Fariha Hasan, Najma Ayub. "Pattern of Drug Resistance and Risk Factors Associated with Development of Drug Resistant Mycobacterium tuberculosis in Pakistan", PLOS ONE, 2016  
Crossref 6 words — < 1%

---

34 scholarhub.ui.ac.id  
Internet 6 words — < 1%

---

EXCLUDE QUOTES ON  
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES OFF  
EXCLUDE MATCHES OFF