

THE EFFECT OF DIABETES SELF MANAGEMENT EDUCATION (DSME) ON SELF EFFICACY IN TYPE II DIABETES MELLITUS PATIENTS

By Subkhan et al

Original Research Article

THE EFFECT OF DIABETES SELF MANAGEMENT EDUCATION (DSME) ON SELF EFFICACY IN TYPE II DIABETES MELLITUS PATIENTS

Mohammad Subkhan¹, Firman Firman^{2*}, Ida Agustiningih^{2,3}

¹ Department of Medicine, Faculty of Medicine, University of Muhammadiyah Surabaya

² Department of Medical Surgical Nursing, Faculty of Medicine, University of Muhammadiyah Surabaya

³ Department of Nursing, Siti Khadijah Hospital, Sepanjang Sidoarjo

*Correspondence:

Firman Firman
Department of Medical Surgical Nursing, Faculty of Medicine, University of Muhammadiyah Surabaya Raya Sutorejo No.59, Surabaya, East Java, 60113
Email: firman@um-surabaya.ac.id

Article Info:

Received: May, 10th, 2024

Revised: June, 10th, 2024

Accepted: June, 10th, 2024

DOI:

<https://doi.org/10.36720/nhjk.v13i1.651>

Abstract

Background: Diabetes mellitus (DM) is a health problem in the world. Currently, the number of DM cases globally has reached 537 million people, and Indonesia ranks fifth highest in the world. Low knowledge and dietary compliance are a problem that continues to contribute to new cases and relapses of old cases characterized by uncontrolled increases in blood sugar. This is caused by low self-efficacy which is characterized by feelings of hopelessness.

Objectives: The aim of this research was to determine the effect of Diabetes Self-Management Education (DSME) on self-efficacy in type II Diabetes Mellitus patients.

Methods: This study used a Quasy-experimental design, with a pretest posttest equivalent control group design, the sample size was 68 type II DM patients, consisting of 34 intervention groups and 34 control groups. Self-efficacy data was collected using DMSES (Diabetes Management Self Efficacy Scale) questionnaire. Univariate analysis was used to describe the frequency distribution of respondents, mean, median and standard deviation. Next, bivariate analysis used the Wilcoxon test. Result: There was a significant difference in the difference in mean value self-efficacy between before and after intervention, with value (p-value 0.000<0.05).

Results: There was a significant difference in the difference in mean value self-efficacy between before and after intervention, with value (p-value 0.000<0.05).

Conclusion: *Diabetes Management Self Efficacy Scale is effective increase self-efficacy. Thus, DSME can be an alternative intervention to overcome problems and prevent complications in DM patients.*

Keywords: *Diabetes Self-Management Education, Diabetes Management Self Efficacy Scale*

INTRODUCTION

Diabetes mellitus is a chronic disease that has an impact on various aspects of life,

both physical, psychological and economic (Boyko et al., 2022). People who suffer from DM (Diabetes mellitus) are characterized by an

uncontrolled increase in blood sugar, which will gradually cause health problems such as retinopathy, which causes the eyes to become blurry and the microvasculature is characterized by tingling. (Malik et al., 2022). This condition becomes worse if you do not adhere to the recommended diet (Page-Reeves et al., 2023). Low knowledge and self-efficacy cause a person to be non-compliant with a diet. In fact, unhealthy eating patterns are the main contributor to the increase in cases of type 2 diabetes mellitus in the world (Han et al., 2022).

According to data from the IDF (International Diabetes Federation), currently the number of cases in the world has reached 537 million people, which is expected to continue to increase by 46%, namely to 643 million people in 2030, and 783 million people in 2045. (IDF, 2021). Indonesia is in fifth place with the highest number of diabetes mellitus cases in the world reaching 19.5 million people, after China is in first place reaching 140.9 million people, then followed by India with 74.2 million people, Pakistan with 33 million people and America with 32.2 million people. The number of cases is expected to continue to increase, along with unhealthy lifestyles. Indonesia is expected to experience an increase of 28.6 million people in 2045 (WHO (World Health Organization), 2021). The same thing also happened at Siti Khadijah Paeng Hospital, where there was an increase in the number of type 2 DM cases in 2023 to 2024, reaching 438, from 2022 which was only 275 people.

The high number of diabetes mellitus cases in Indonesia is influenced by several factors, such as increased body weight or obesity (29.0%). Obesity is caused by poor eating patterns (77.4%), continuous excess nutritional intake which causes excess fat stores in the body. This fat tissue is an active endocrine tissue that releases adipose cytokines. Cytokines will interfere with the insulin signaling pathway, causing insulin resistance. Another factor, namely lack of

activity (64.5%), not doing physical activity, has not been proven to increase the risk of developing type 2 DM, this is probably caused by physical activity carried out daily (such as walking to the market, hoeing, washing, gardening) not including doing activities. Physique (Dungga et al., 2024).

Confidence in the abilities of people with diabetes mellitus to be more compliant with diet and self-management in order to recover from their illness. A previous study explained that preventing complications in diabetes mellitus is by controlling blood sugar (Malik et al., 2022). Well-controlled blood sugar can reduce the risk of retinopathy (Petersen et al., 2022 ; Wan et al., 2022). Higher self-efficacy indicates good self-management, and being more compliant with treatment and tending to have a healthy lifestyle. The results of a randomized study show that there is a relationship between self-efficacy and a decrease in HBA1C (Handelsman et al., 2019). Peer learning or fellow DM patients is effective in improving better self-management (Seuring et al., 2019). The results of the review study explain that Diabetes self-management education (DSME) is effective in increasing the knowledge and behavior of diabetes mellitus patients (Camargo-Plazas et al., 2023). DSME can increase knowledge and self-care behavior (Roberta et al., 2023).

Based on several studies, improving a patient's knowledge and behavior is very important, to encourage better health, so that complications do not occur. Apart from that, previous research has not specifically explained how to increase self-efficacy and adherence to diet. Diabetes self-management education (DSME) is an intervention that provides health education about DM, education about diet, physical activity and health monitoring. (He et al., 2023).

Objective(s): The aim of this research is to determine the effectiveness of the Diabetes self-management education (DSME)

intervention on self-efficacy in type II DM patients.

6 METHODS

Study Design

This research used a quasy experimental design, with a pretest posttest equivalent control group design with repeated measurements.

Research Subject

Participants were recruited using convenience sampling, sampling was carried out at Siti Khadijah Hospital, Sepanjang Sidoarjo, East Java, Indonesia, from February to March 2024. Respondents met the following inclusion criteria; Type II DM patients, patients visiting the internal medicine clinic, cooperative, willing to be research samples. And were excluded if their consciousness decreased, they had mental disorders and they refused to participate in this research. The total sample was 68 people, consisting of an intervention group and a control group, each group totaling 34 people.

Instruments

The first measuring tool used in this research³⁴ is collecting sociodemographic data using a questionnaire including (age, gender, education level, marital status, employment, income and¹² comorbidities). To measure self-efficacy, the DMSES (Diabetes Management Self-Efficacy Scale) questionnaire was used. Providing Diabetes Self-Management Education treatment uses the SPO Diabetes Self-Management Education (DSME) check list.

Intervention

The Diabetes Self-Management Education (DSME) intervention procedure is carried out three times every day, in the morning, after have lunch and the afternoon. which consists of three sessions, the first session provides material about diabetes

mellitus, the second session was about complications and their prevention, and the third session provides reinforcement, motivation and support. Each session lasts 15-20 minutes for 7 days. Pre-test given the intervention, self-efficacy¹⁴ and sociodemographic were measured in the intervention group and control group.¹⁴ furthermore, DSME intervention was given to the intervention group, and the control group was only given standard intervention from the hospital. The intervention stage, starting from the preparation stage, begins with giving explain to the respondent the procedures to be carried out, maintain the respondent's privacy and create a comfortable atmosphere. Implementation stage ask the patient to lie in a supine or sitting position and relax, furthermore providing intervention with Diabetes Self-Management Education, the next is the termination or evaluation stage, and gives them the opportunity to provide feedback on the therapy that has been carried out.

19 Data Analysis

Data were analyzed using SPSS software (Version 25.0; Chicago IL USA), to describe data on gender, education level and marital status, self-efficacy and dietary compliance, frequency and percentage distributions were used. Meanwhile, age and blood sugar values are presented in the form of mean, median, min-max and standard deviation. For further analysis, to determine the difference in the¹³ mean pretest and posttest values because the data was not normally distributed, the Wilcoxon test was carried out with a significant value (p-value < 0.005).

Ethical Consideration

The procedures in this study involving human participants by administering the DSME²³ intervention have been approved and declared in accordance with the ethical standards of the Health Research Ethics Committee (KEPK) of the Siti Khadijah Hospital, Sepanjang Sidoarjo,

with registration number (No.018/KET-KEPK/6-2023). All participants in this study have received an explanation and expressed their willingness to sign the informed consent that was given.

RESULTS

Univariate

Table 1. Characteristics of Respondents

Based on Age					
	Group	Mean	Min-Max	Median	Std. Dev
Age	Intervention	53.38	43-67	53.90	6,344
	Control	55.94	46-65	54.50	5,908

Based on the table above, it showed that the characteristics of respondents according to the average age of respondents in the intervention group was 53.38 years old and the median value was 53.90 with standard deviation 6,344. The youngest is 43 years old and the oldest is 67 years old. Meanwhile, the average age in the control group was 55.94 years and the median value was 54.50 with standard deviation 5,908. The youngest is 46 years old and the oldest is 65 years old.

Table 2. Characteristics of Respondents

Variable	Intervention Group		Control group	
	n	%	n	%
Gender				
Man	11	32.4	15	44.2
Woman	23	67.6	19	55.8
Total	34	100	34	100
Level of education				
Elementary school	4	11.8	3	8.8
Junior High School	9	26.5	8	23.5
Senior High School	11	32.4	11	32.4
Diploma/Bachelor's degree	10	29.4	12	35.3
Total	34	100	34	100
Marital status				
Marry	27	79.4	23	67.6
Widow widower	7	20.6	11	32.4
Total	34	100	34	100

Job

Doesn't work	7	20.6	12	35.2
Work	23	67.6	20	59.8
Pension	4	11.8	2	5.0
Total	34	100	34	100

Income

< Rp. 1,000,000	5	14.8	9	26.4
Rp. 1,000,000 - Rp. 2,000,000	3	8.9	7	20.6
> Rp. 2,000,000 - Rp. 3,000,000	7	20.5	6	17.7
> Rp. 3,000,000 - Rp. 4,000,000	10	29.4	4	11.7
>4,000,000	9	26.4	8	23.6
Total	34	100	34	100

Comorbid

Hypertension	28	83.3	24	71.4
Heart disease	2	5.0	8	23.6
CKD	4	11.7	2	5.0
Total	34	100	34	100

The table above showed that the distribution of respondents based on gender characteristics in the intervention group, most of the respondents were women, 23 people (67.6%) and in the control group, the majority of respondents were also women, 19 people (55.8%). According to the educational level of respondents in the intervention group, the majority were high school, 11 people (32.4%), while in the control group, the majority were Diploma/Bachelor's degrees, 12 people (35.3%). The marital status of most respondents was married, in the intervention group 27 people (79.4%) and in the control group 23 people (67.6%). Most of the respondents were working, the intervention group was 23 people (67.6%). Meanwhile, 20 people (59.8%) worked in the control group. The income level the intervention group was mostly income > Rp. 3,000,000 - Rp. 4,000,000, as many as 10 people (29.4%). Meanwhile, most of the control group had income < Rp. 1,000,000, as many as 9 people (26.4%). Most of the comorbid diseases experienced were hypertension, 28

36 people in the intervention group (83.3%), and 24 people in the control group (71.4%).

Table 3 Distribution Overview *self-efficacy* Pre-test and Post-test Results

Intervention Group		Pre-test		Post-test	
Variable	n	%	n	%	
Tall	9	26.5	31	91.2	
Low	25	73.5	3	8.8	
Total	34	100.0	34	100.0	
Control group		Pre-test		Post-test	
Variable	n	%	n	%	
Tall	12	35.3	10	29.4	
Low	22	64.7	24	70.6	
Total	34	100.0	34	100.0	

Based on the table above distribution *self-efficacy* Most of the respondents in the Pre-test intervention group were given the Diabetes Self-Management Education (DSME) intervention *self-efficacy* low by 25 people (73.5%). And the post-test was given the most intervention *self-efficacy* high as many as 31 people (91.2%). Meanwhile distribution *self-efficacy* Most of the respondents in the control group had pre-test results *self-efficacy* low as many as 22 people (64.7%), and post-test most of the distribution *self-efficacy* low amounting to 24 people (70.6%).

Bivariate

Table 4. Mean Difference Analysis *Self-Efficacy* Pre-test and Post-test Results

<i>Self-efficacy</i>	Different Means	P-Value
Intervention Group		
Pre-test - Post-test	11.50	0,000*
Control Group		
Pre-test - Post-test	2.50	0.317

*Significant at $\alpha < 0.05$, by Wilcoxon test

From the results of the Wilcoxon analysis above, it showed that there was a significant influence on the mean *self-efficacy* of the intervention group. Pre-test and post-test were given the intervention *Diabetes Self-Management Education* with value (P-Value $0.000 < 0.05$). Thus, this shows that there are significant differences in the mean values *self-efficacy* in the intervention group, where the difference in change from the pre-test and corresponding was 11.5. Meanwhile, the results of the analysis in the control group showed that there was no significant influence between the mean value of *self-efficacy* from the pre-test and post-test results, as evidenced by the value (p-value $0.317 > 0.05$).

DISCUSSION

The findings of this study showed that there was a significant influence on the mean *self-efficacy* of the intervention group. Pre-test and post-test were given the intervention *Diabetes Self-Management Education* with value (p-value $0.000 < 0.05$) indicates that there was a significant difference in the mean value *self-efficacy* in the intervention group. Meanwhile, the results of the analysis in the control group showed that there was no significant influence between the mean value of *self-efficacy* from the pre-test and post-test results, as evidenced by the value (p-value $0.317 > 0.05$). Based on these results, the intervention group had an influence on increasing *self-efficacy* scores.

Self-efficacy is the belief that a person can carry out a certain behavior to achieve the desired goal because high *self-efficacy* can be influenced by several factors such as *Diabetes Self-Management Education*. The results of this research are in line with previous research conducted by Rahmawati, (2016), because based on this research it is not only the knowledge of individuals, families and communities in efforts to control type 2 DM but also the use of health information, changes in attitudes, changes in behavior and increased

compliance with treatment program and improving the quality of life of type 2 DM sufferers.

DSME can also overcome distress in type 2 DM patients ($p=0.001$), because diabetes distress refers to negative psychological reactions related to emotional stress and worry about the situation, so the effect of giving DSME to DM patients who face complex chronic diseases and stress in reducing the burden of diabetes can worsen the condition after research showed that there was an effect of giving DSME/S on reducing diabetes distress (Nurkamilah & Widayati, 2018). Clients who are facilitated with knowledge and skills in carrying out self-care for DM through providing DSME/S education, which is support-based health education in a directed and structured manner, can learn how to solve problems experienced by DM patients. Diabetes Self-Management Education increases patient knowledge about independent self-care compared to simply providing knowledge about diabetes (Murtaqib., 2019).

Diabetes Self-Management Education (DSME) is a strategy for transferring knowledge as well as raising patient enthusiasm, so that they will demonstrate healthy behavior. Diabetes is a chronic disease that requires lifelong treatment. Therefore, comprehensive treatment is required. Inappropriate care has broad implications for patients and families (Murtiningsih et al., 2021). This is because the DSME intervention group consistently adheres to the four pillars of diabetes, namely education, nutritional management, implementation of physical activity, and daily drug therapy. (Malik et al., 2022). Different from the control group intervention. In the control group intervention, the other three pillars of diabetes management were not implemented, and education was only provided about the general description of diabetes, without special monitoring of diet management or physical activity. So that the DSME intervention group experienced a

6crease in blood sugar levels at any time and the control group experienced an increase in blood sugar levels at any time.

This finding is supported by research conducted by Uly, (2018), showing that there was a significant change in the mean blood glucose levels of respondents in both groups after implementing the intervention. According to Perkeni (2015), there are four main pillars in the treatment of type 2 diabetes patients: education, nutritional therapy, exercise, and pharmacology. Education plays an important role in the treatment of type 2 DM because educating diabetes patients can change their diabetes self-care behavior. According to the researchers' assumptions in this study, DSME interventions rely more on knowledge, nutritional management, physical activity, foot care, stress management, and medication use. They were compared with a control group who were only given information about non-communicable diseases. The results of this study showed that there was a significant difference in HBA1C levels between the treatment group and the control group (Handelsman et al., 2019).

CONCLUSION

There is a significant difference in the mean value self-efficacy between before and after intervention, with value ($P\text{-Value } 0.000 < 0.05$). This shows that *Diabetes Management Self Efficacy Scale* is effective increase self-efficacy. DSME is a form of intervention in health education for diabetes mellitus patients. Health education not only transfers knowledge to patients, but can also increase enthusiasm and confidence in each patient's ability to recover from the DM disease they are experiencing. Thus, DSME can be an alternative intervention to overcome problems in DM patients and prevent complications.

SUGGESTIONS

This study has several limitations, firstly, participants were not recruited

randomly, this is because the number of hospitalized patients was not sufficient according to the specified number, so with these limitations the sample was determined using convenience sampling, meaning the researcher took patients who met the inclusion criteria and willing to be a participant in this research. Therefore, further research needs to be carried out by identifying and controlling possible confounding variables, including drug therapy given to participants during the research.

4 ACKNOWLEDGMENT

The researchers would like to thank the Diktilitbang Council (Higher Education Research and Development Council of Muhammadiyah), which has provided funding assistance for this research, LPPM Muhammadiyah University Surabaya, which has supported this research, and to Siti Khadijah Hospital, Sepanjang Sidoarjo which has grant research permission. Lastly, thank you also to the participants involved in this research.

25 DECLARATION OF CONFLICTING INTEREST

No conflict of interest

FUNDING

The study was funded by Diktilitbang (Higher Education Research and Development Council of Muhammadiyah).

AUTHOR CONTRIBUTION

Mohammad Subkhan: Conduct the study and collect the data.

Firman Firman: Conduct the study, analysis, translation and submit this manuscript.

Ida Agustinarsih: Conduct the study, provided access and coordinated participants.

ORCID

Mohammad Subkhan:

None

Firman Firman:

None

Ida Agustinarsih:

None

REFERENCES

- Boyko, E.J., Zelnick, L.R., Braffett, B.H., Pop-Busui, R., Cowie, C.C., Lorenzi, G.M., Gubitosi-Klug, R., Zinman, B., & de Boer, I.H. (2022). Risk of Foot Ulcer and Lower-Extremity Amputation Among Participants in the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications Study. *Diabetes Care*, 45(2), 357–364. <https://doi.org/10.2337/dc21-1816>
- Camargo-Plazas, P., Robertson, M., Alvarado, B., Paré, G.C., Costa, I.G., & Duhn, L. (2023). Diabetes self-management education (DSME) for older persons in Western countries: A scoping review. In *PLoS ONE* (Vol. 18, Issue 8 August). <https://doi.org/10.1371/journal.pone.0288797>
- Han, H., Cao, Y., Feng, C., Zheng, Y., Dhana, K., Zhu, S., Shang, C., Yuan, C., & Zong, G. (2022). Association of a Healthy Lifestyle With All-Cause and Cause-Specific Mortality Among Individuals With Type 2 Diabetes: A Prospective Study in UK Biobank. *Diabetes Care*, 45(2), 319–329. <https://doi.org/10.2337/dc21-1512>
- Handelsman, Y., Chovanes, C., Dex, T., Giorgino, F., Skolnik, N., Souhami, E., Stager, W., Niemoeller, E., & Frias, J.P. (2019). Efficacy and safety of insulin glargine/lixisenatide (iGlarLixi) fixed-ratio combination in older adults with type 2 diabetes. *Journal of Diabetes and Its Complications*, 33(3), 236–242.

- <https://doi.org/10.1016/j.jdiacom.2018.11.009>
- He, Q., Liu, Y., Lin, K., Zhao, F., Guo, H., & Shen, Y. (2023). Diabetes self-management and its related factors among Chinese young adults with type 2 diabetes mellitus. *Nursing Open*, 10(9), 6125–6135. <https://doi.org/10.1002/nop2.1834>
- IDF. (2021). International Diabetes Federation. In *Diabetes Research and Clinical Practice* (Vol. 102, Issue 2). <https://doi.org/10.1016/j.diabres.2013.10.013>
- Dungga EF, & Indiyanti, Y., (2024). Risk Factors For Type 2 Diabetes Mellitus Patients At The Monano Health Center, North Gorontalo District. *Medicine, F., Gorontalo.*, 6(1), 40–56.
- Nurkamilah, N. and Widayati, N. (2018) '(DSME/S) on Diabetes Distress in Type 2 Diabetes Mellitus Patients at RSD dr. Soebandi Jember [DSME/S] on Diabetes Distress in Patients with Type 2 Diabetes Mellitus in dr. Soebandi Hospital of Jember)', *e-Jurnal of Health Literature*, 6(1), pp. 133–140.
- Malik, F.S., Sauder, KA, Isom, S., Reboussin, B.A., Dabelea, D., Lawrence, J.M., Roberts, A., Mayer-Davis, E.J., Marcovina, S., Dolan, L., Igudesman, D., & Pihoker, C. (2022). Trends in Glycemic Control Among Youth and Young Adults With Diabetes: The SEARCH for Diabetes in Youth Study. *Diabetes Care*, 45(2), 285–294. <https://doi.org/10.2337/dc21-0507>
- Murtaqib (2019) 'The Effect of Self Management Education and Support in Improving the Self-Efficacy of Patients with Type 2 Diabetes Mellitus', *JNC*, 2(1), pp. 1–9.
- Murtiningsih, MK, Pandelaki, K., & Sedli, BP (2021). Lifestyle as a Risk Factor for Type 2 Diabetes Mellitus. *E-CliniC*.
- Page-Reeves, J., Murray-Krezan, C., Burge, M.R., Mishra, S.I., Regino, L., Bleecker, M., Perez, D., McGrew, H.C., Bearer, E.L., & Erhardt, E. (2023). A patient-centered comparative effectiveness research study of culturally appropriate options for diabetes self-management. *MedRxiv : The Preprint Server for Health Sciences*. <https://doi.org/10.1101/2023.01.31.23285236>
- PERKENI (2015). Management and Prevention of Type 2 Diabetes Mellitus in Indonesia, PERKENI, Jakarta: 13
- Petersen, G. B., Byberg, S., Vistisen, D., Fangel, M. V., Vorum, H., Joensen, L. E., & Kristensen, J. K. (2022). Factors Associated With Nonattendance in a Nationwide Screening Program for Diabetic Retinopathy: A Register-Based Cohort Study. *Diabetes Care*, 45(2), 303–310. <https://doi.org/10.2337/dc21-1380>
- Rahmawati (2016) 'The Effect of the Diabetes Self-Management Education Program on Self-Management in Patients with Diabetes Mellitus Type 2 Effects of Diabetes Self-Management Education Program on Self-Management in Patients with Diabetes Mellitus Type 2', *Journal of Nursing Science*, 4:1, pp. 46–58
- Roberta, L., Amoakoh-Coleman, M., Djobalar, B., Grobbee, D.E., Adjei, G.O., & Klipstein-Grobusch, K. (2023). Diabetes self-management education interventions and self-management in low resource settings; a mixed methods study. *PLOS ONE*, 18(7 July), 1–19. <https://doi.org/10.1371/journal.pone.0286974>
- Seuring, T., Marthoenis, Rhode, S., Rogge, L., Rau, H., Besançon, S., Zufry, H., Sofyan, H., & Vollmer, S. (2019). Using peer education to improve diabetes management and outcomes in a low-income setting: A randomized controlled trial. *Trials*, 20(1), 1–8. <https://doi.org/10.1186/s13063-019-3656-1>

Uly Agustine, LRRW (2018). Factors that Influence the Level of Adherence to Taking Medication in Diabetes Mellitus Sufferers Who Seek Treatment at the A and A Rahmat Waingapu Love Service Foundation Medical Center. *Journal of Primary Health Care*, 3(1). <https://doi.org/Journal> DOI: <https://doi.org/10.31965/jkp>

Wan, Y., Tube, FK, Lee, DH, Fung, TT, Willett, WC, & Giovannucci, EL (2022). Dietary Insulinemic Potential and Risk of

Total and Cause-Specific Mortality in the Nurses' Health Study and the Health Professionals Follow-up Study. *Diabetes Care*, 45(2), 451–459. <https://doi.org/10.2337/dc21-1530>

WHO (World Health Organization). (2021). Advisory Group on Diabetes. December, 1–21.

Cite this article as: Subkhan, M., et al. (2024). The Effect of Diabetes Self-Management Education (DSME) on Self Efficacy in Type II Diabetes Mellitus Patients. *Nurse and Health: Jurnal Keperawatan*, 13 (1), 127-135. <https://doi.org/10.36720/nhjk.v13i1.651>

THE EFFECT OF DIABETES SELF MANAGEMENT EDUCATION (DSME) ON SELF EFFICACY IN TYPE II DIABETES MELLITUS PATIENTS

ORIGINALITY REPORT

24%

SIMILARITY INDEX

PRIMARY SOURCES

- 1** repository.unej.ac.id 55 words — 2%
Internet
- 2** Surya Mustika Sari, Henny Vidia Effendy. "THE EFFECT OF HEALTH EDUCATION ABOUT MENARCHE ON ANXIETY IN FACING MENARCHE IN 5TH AND 6TH GRADE STUDENTS", Nurse and Health: Jurnal Keperawatan, 2019 50 words — 1%
Crossref
- 3** Meri Afridayani, Yohana Ika Prastiwi, Khudazi Aulawi, Ibrahim Rahmat, Hera Nirwati, Haryani Haryani. "Relationship between hand hygiene behavior and Staphylococcus aureus colonization on cell phones of nurses in the intensive care unit", Belitung Nursing Journal, 2021 42 words — 1%
Crossref
- 4** tmfv.com.ua 39 words — 1%
Internet
- 5** jurnal.syntax-idea.co.id 38 words — 1%
Internet
- 6** Effatul Afifah, Detty Siti Nurdiati, Hamam Hadi, Zainal Muttaqien Sofro, Ahmad Hamim Sadewa, Arif Sabta Aji. "The effect of social Nervous (SaSo) exercise with 36 words — 1%

insulin resistance (HOMA-IR) on gestational diabetes mellitus among Indonesian pregnant women: a quasi-experimental study", Research Square Platform LLC, 2021

Crossref Posted Content

7	jurnal.unimus.ac.id Internet	30 words — 1%
8	ibdigital.uib.es Internet	29 words — 1%
9	banrjournal.org Internet	27 words — 1%
10	jurnal.umpp.ac.id Internet	22 words — 1%
11	findresearcher.sdu.dk Internet	21 words — 1%
12	www.coursehero.com Internet	18 words — 1%
13	jurnal.ikta.ac.id Internet	16 words — < 1%
14	publications.aston.ac.uk Internet	16 words — < 1%
15	www.ncbi.nlm.nih.gov Internet	16 words — < 1%
16	Marwan Rosyadi, Atsushi Umemura, Rion Takahashi, Junji Tamura, Takamori Uchiumi, Masanori Mori, Hiroki Miyata. "A simplified model design of MMC-HVDC transmission system for steady state and transient	15 words — < 1%

stability analyses", International Journal of Power Electronics and Drive Systems (IJPEDS), 2023

Crossref

17 Sri Zuliarni, Dwi Kartikasari, Bambang Hendrawan, Siti Sri Windrayati Siregar. "The impact of buying intention of global fashion on local substitute: The role of product design and price", Heliyon, 2023

Crossref

14 words — < 1%

18 repository.ub.ac.id

Internet

14 words — < 1%

19 www.mdpi.com

Internet

14 words — < 1%

20 Mann, Jim, Truswell, Stewart, Hodson, Leanne. "Essentials of Human Nutrition 6e", Essentials of Human Nutrition 6e, 2023

Publications

13 words — < 1%

21 Straits, Bruce C.. "Social Research", Oxford University Press

Publications

13 words — < 1%

22 eprints.ulm.ac.id

Internet

13 words — < 1%

23 link.springer.com

Internet

13 words — < 1%

24 Indriyani Indriyani, Mellyana Cahyadi. "The Relationship of Obesity with Range Of Motion (ROM) On The Ankle", Biomedical Journal of Indonesia, 2023

Crossref

12 words — < 1%

25 Salsabilla Rohadatul 'Aisy Sunaryo, Azam David Saifullah, Sri Mulyani. "KNOWLEDGE AND

12 words — < 1%

ATTITUDES TOWARD PEOPLE WITH DEMENTIA AMONG
NURSING STUDENTS IN YOGYAKARTA, INDONESIA", Belitung
Nursing Journal, 2020

Crossref

-
- 26 ejournal.iainpalopo.ac.id 12 words — < 1%
Internet
-
- 27 goodwoodpub.com 12 words — < 1%
Internet
-
- 28 repository.sustech.edu 12 words — < 1%
Internet
-
- 29 Agus Purnama, Mua Melawati. "The Benefits of Dates Ruthab on Blood Volume On Labor Stage 4th", Journal Of Nursing Practice, 2019 11 words — < 1%
Crossref
-
- 30 Kusuma Estu Werdani, Izzatul Arifah, Anggi Putri Aria Gita, Mislailatul Kholifah Rahma et al. "Contraception used among women of childbearing age during the pandemic COVID-19", International Journal of Public Health Science (IJPHS), 2024 11 words — < 1%
Crossref
-
- 31 Raju Kapadia, Mita Agustina, Niya Fitarsih. "Implementation of Online Problem-Based Learning And Nursing Students' Engagement With Modified Online Discussion With Seven Jumps (MOODS7)", Jurnal Keperawatan Terpadu (Integrated Nursing Journal), 2023 11 words — < 1%
Crossref
-
- 32 Titiek Idayanti, Kurnia Indriyanti, Widya Anggraeni. "THE EFFECT OF EARLY BREASTFEEDING INITIATION (IMD) ON NEW BORN BABY TEMPERATURE IN BPS HEPPY RINA, S.ST, SEDURI VILLAGE-

MOJOSARI AND BPS FIFIT, S.ST, PANJER VILLAGE-MOJOSARI",
Nurse and Health: Jurnal Keperawatan, 2019

Crossref

33	repository.seafdec.org Internet	11 words — < 1%
34	www.ephysician.ir Internet	11 words — < 1%
35	diabetesjournals.org Internet	10 words — < 1%
36	e-journal.unair.ac.id Internet	10 words — < 1%
37	journals.plos.org Internet	10 words — < 1%
38	jurnal.feb-umi.id Internet	10 words — < 1%
39	www.jcpsp.pk Internet	10 words — < 1%
40	Enza Gucciardi, Margaret DeMelo, Ruth N. Lee, Sherry L. Grace. "Assessment of two culturally competent Diabetes education methods: Individual versus Individual plus Group education in Canadian Portuguese adults with Type 2 Diabetes", Ethnicity & Health, 2007 Crossref	9 words — < 1%
41	ejournal.delihusada.ac.id Internet	9 words — < 1%
42	ejournal.seaninstitute.or.id Internet	9 words — < 1%

- 43 ejurnal.ung.ac.id
Internet 9 words — < 1%
-
- 44 sloap.org
Internet 9 words — < 1%
-
- 45 David M. Maahs, Priya Prahalad, Darja Šmigoc Schweiger, Shlomit Shalitin. "Diabetes Technology and Therapy in the Pediatric Age Group", *Diabetes Technology & Therapeutics*, 2023
Crossref 8 words — < 1%
-
- 46 Lono Wijayanti. "ROLE OF FAMILY IN IMPLEMENTATION OF DIABETES EXERCISE IN TYPE 2 DIABETES MELITUS PATIENTS", *Nurse and Health: Jurnal Keperawatan*, 2019
Crossref 8 words — < 1%
-
- 47 Rondhianto Rondhianto, Akhmad Zainur Ridla, Fandi Ahmad Kurniawan Setia Budi. "Analysis of Psychosocial Factors Affecting Dietary Behavior among People with Type 2 Diabetes Mellitus in Indonesia", *Jurnal Keperawatan Soedirman*, 2023
Crossref 8 words — < 1%
-
- 48 assets-eu.researchsquare.com
Internet 8 words — < 1%
-
- 49 brieflands.com
Internet 8 words — < 1%
-
- 50 garuda.ristekbrin.go.id
Internet 8 words — < 1%
-
- 51 talenta.usu.ac.id
Internet 8 words — < 1%

52 turkjphysiotherrehabil.org 8 words — < 1%
Internet

53 www.heartindiabetes.com 8 words — < 1%
Internet

54 Eni Sumarliyah, Dede Nasrullah, Sukadiono ., 6 words — < 1%
Desta Pankyano. "Lowering blood sugar levels with
kersen leaf stew (muntingia calabura l) and black onion (alium
sativum) extract in mice (mus musculus)", International Journal
of Advanced Nursing Studies, 2020
Crossref

55 Qiao He, Yu Liu, Keke Lin, Fang Zhao, Hong Guo, 6 words — < 1%
Yina Shen. "Diabetes self-management and its
related factors among Chinese young adults with type 2
diabetes mellitus", Nursing Open, 2023
Crossref

56 pure.rug.nl 6 words — < 1%
Internet

EXCLUDE QUOTES ON

EXCLUDE SOURCES OFF

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF