EFFECTIVENESS OF WHATSAPP BOT-BASED EDUCATIONAL MEDIA AND LECTURES ON INCREASING MOTHER'S KNOWLEDGE ABOUT GASTROENTERITIS IN CHILDREN

By Ar Royo et al

Ar Royo, S. P., et al. (2024). Nurse and Health: Jurnal Keperawatan. 13 (2): 19-29 http://ejournal-kertacendekia.id/index.php/nhjk/index

Original Research Article

EFFECTIVENESS OF WHATSAPP BOT-BASED EDUCATIONAL MEDIA AND LECTURES ON INCREASING MOTHER'S KNOWLEDGE ABOUT GASTROENTERITIS IN CHILDREN

Sonia Puspita Ar Royo¹, Anis Ika Nur Rohmah^{1*}, Ika Rizki Anggraini¹, Nur Melliza¹

¹University of Muhammadiyah Malang Malang, Indonesia

*Correspondence:



University of Muhammadiyah Malang Malang

Raya Tlogomas No. 246, Malang, East

Java, Indonesia 65144 66 Email: anisikanur@umm.ac.id

Article Info:

Received: June, 25th, 2024 Revised: Sember, 18th, 2024 Accepted: September, 26th, 2024

DOI:

https://doi.org/10.36720/nhjk.v13i2.663

Abstract

Background: Gastroenteritis is the main cause of death in children under five years of age with symptoms of dehydration, which causes high mortality rates. Gastroenteritis risk factors include poor hygiene, unclean water supplies, and lack of parental knowledge. Providing health education regarding gastroenteritis is very important a increase mother's knowledge. WhatsApp bot and lectures are effective media that can be used in health education.

Objectives: To determine the effectiveness of WhatsApp bot-based educational media and lectures in increasing knowledge of Gastroenteritis among hou

Methods: The research method used was quasi-experimental using a two-group design posttest only. The research was conducted in Singosari, Malang, using two different groups with a sample size of 50. Group 01 was given education using WhatsApp bot media, while group 02 was given education using lecture media. The research data was tested using the Mann-Whitney test in SPSS version 25.

Results: The demographic data obtained shows that the average number of respondents is (60%) not working or a housewife. The research results show a difference between giving education using the lecture method and WhatsApp bot media on housewives' knowledge level. Mother's knowledge about gastroenteritis increased after being given education using two different media. The group using lecture media had a better increase in knowledge than the group using WhatsApp bot media.

Conclusion: Health education regarding gastroenteritis based on lecture media is more effective than WhatsApp bot media in increasing mothers' knowledge about preventing gastroenteritis in children.

Keywords: Gastroenteritis, Knowledge, Lectures, WhatsApp bot

© 2024 The Authors. Nurse and Health: Nursing Journal Published by Institute for Research and Community Service - Health Polytechnic of Kerta Scholar

This is an Open Access Article distributed under the terms of the Creative Commons Attribution – Non-Commercial 4.0 (CC BY-NC) 4.0) which allows others to remix, tweak, and build upon the work non-commercially as long as the original work is properly cited. The new creations are not necessarily licensed under identical terms.

E-ISSN 2623-2448 P-ISSN 2088-9909

INTRODUCTION

Gastroenteritis is inflammation that occurs in the stomach and results in changes in

vital signs (pulse, blood pressure, heart rate) leading to frequent diarrhea and nausea, vomiting (Doris, 2021; Qori Nurul Isnaini,

2021). Gastroenteritis is caused by many factors, including a lack of nutritional intake, a lack of environmental health, and an unclean style (Yona & Iriani, 2022). Gastroenteritis is the main leading of death in children under five years of age (toddlers) which results in dehydration, triggering high morbidity, and mortality rates (Arsi, Afdhal, & Fatrida, 2021; Khoiriyah, Widjajanegara, & Sjafei, 2021; Nari, 2019; Rachman & Nurjanah, 2019). The World Health Organization (WHO), (2018) reported that gastroenteritis attacks 66 million people in the world with a death rate of around 546,000 in children under five each year, while in Indonesia, the Indonesian Ministry of Health reported that in 2017 gastroenteritis was recorded as occurring 21 times spread across 12 provinces and 17 districts/cities in Indonesia with a total of 1,725 sufferers. Gastroenteritis has also been reported to cause seizures that require hospitalization (Heydarian et al., 2019). In 2021, the prevalence of gastroenteritis in Malang City in the toddler group is 14.1% and in all age groups is 14.8% (Malang City Health Service, 2022). Aditya's research in (Herlina Alvianti Ningsih, Yulia Wardita, 2021) in Malang Regency, Singosari District, it was revealed that 21 babies aged 0-6 months experienced gastroenteritis, as well as phenomena that exist in the Singosari area such as a lack of health education regarding gastroenteritis and a lack of parental knowledge regarding gastroenteritis.

Gastroenteritis can occur because food is not processed properly and the water is contaminated (Rahmawati & Arifiyanti, 2020). There is a relationship between parental practices in providing food and the incidence of gastroenteritis (Siahaan, Aritonang, & Ashar, 2020). In addition, the quality of clean water also has a significant relationship with the incidence of gastroenteritis in toddlers (p<0.05)(Rimbawati & Surahman, 2019). There is a relationship between these phenomena, as a Muslim it is very contrary to the word of Allah in the QS Al-Baqarah [2]: 222).

ٱلْمُتَطَهَرِينِ وَيُحِبُّ ٱلتَّوَّٰبِينَ يُحِبُّ ٱللَّهَ إِنَّ.

".....Indeed, Allah loves those who repent and those who purify themselves" (QS. Al-Baqarah [2]: 222).

In these words, a Muslim should be able to maintain personal hygiene regarding the environment and food. In this case, there is a relationship between the level of education and a person's hygiene (p<0.001) (Ariani, 2022).

The high incidence of gastroenteritis cannot be arrated from the role of parents, especially the role of mothers in preventing and treating diseases in children (Komara et al., 2020). The mother's knowledge gastroenteritis shows that mothers know everything related to the disease so that it can reduce the incidence of gastroenteritis in children (Arianti & Jaya, 2022; Purnamasari & Putri, 2023). Prevention of gastroenteritis through increasing maternal knowledge can be achieved by providing health education through the media (Febiyana, Saputri, Sari, & Sari, 2024). Effective media that can be used are lectures and WhatsApp bots because they provide useful information and services (Dewi, 2022).

Lectures are often used in health education to disseminate information and discuss each other but are also considered boring media (Mutmawardina et al., 2023; Susilowati & Suwarni, 2023). WhatsApp is used in mobile health interventions that utilize automatic chat features to send information (Manji et al., 2021). WhatsApp as a two-way communication platform serves to deliver education and provide users with learning opportunities at any time (Giansanti, 2020; Salam et al., 2021). Chatbot in WhatsApp is a chat system that makes providing information easier and does not limit user consultation (Umma et al., 2024). Several studies have conducted research using lecture WhatsApp bot methods, but no researchers have compared lectures and WhatsApp bot.

Based on the explanation above, the author initiated a health management innovation by providing education based on

lecture methods and WhatsApp bot media on the number of gastroenteritis sufferers in children and toddlers 63

Objective: This research aims to determine the effectiveness of educational media based on WhatsApp bots and lectures in increasing knowledge of Gastroenteritis among housewives.

METHODS

Study Design

The research method was quasiexperimental with two groups posttest-only design. The activity was carried out by providing education using WhatsApp bot media in group 01 and providing education using the lecture method in group 02 at different times.

Settings

The research was conducted on two groups in Singosari District.

Reseggh Subject

The sampling technique used total sampling. Therefore, the sample in this study was the entire population, namely 50 people. In each group, the determination of each sample was not carried out randomly so that it was by the existing population, while the division between the WhatsApp bot experimental and lecture groups was carried out randomly using a random draw. The population taken in this study were all mothers in Group 01 and Group 02 Singosari District, especially mothers who already had children and toddlers with a total sample of 50 mothers.

Instruments

The data collected in this research is primary data collected directly from the data source. The independent variable in the st source. The independent variable in the st source was gastroenteritis education, while the dependent variable was the level of knowledge. The instrument used was the Mother's Knowledge L Questionnaire about Gastroenteritis which had been tested for

validity and reliability by previous researchers.

The statistical test used to analyze the correlation between variables is the Mann-Whitney.

Intervention

This research used two experimental groups, there was no control group and only used posttest in two groups. The intervention was given using different media for each group to compare the two media used. The activity was carried out by starting with providing education using WhatsApp bot media to group 01. Researchers used WhatsApp bot to deliver the material and respondents could immediately try using the WhatsApp bot application. On WhatsApp bot there is an explanation about gastroenteritis starting from the definition, signs, symptoms, causes, risk factors, prevention, and treatment. By saying the keyword "hello mother", WhatsApp will respond to several question codes, after which you can choose a code and then write the appropriate code. WhatsApp bot will provide 10 different codes such as code #01 which means the definition of gastroenteritis so that when responders type #01 a detailed explanation of gastroenteritis will appear along with valid references from the Ministry of Health and the way the other codes work is also the same. Questions on WhatsApp bot are unlimited and users can consult 24 hours a day. In the WhatsApp bot group, the intervention was given twice at different times with a duration of 2 hours.

In the intervention in group 02, education was also provided using the lecture method. After the education about gastroenteritis was completed, a question-answer session was held. In group 02, researchers used the lecture method where the material was delivered orally so that respondents could listen carefully to the material given and the material in the lecture group was also the same as the material in the WhatsApp Bot group. In the lecture group, intervention was given twice at different times

with a duration of 2 hours. The intervention time between group 01 and group 02 is different.

Data Analysis

The measurement process used Statistical Package for Social Science (SPSS) version 25 software.

Ethical Considerations

This research 23 received approval and permission from the Faculty of Health Sciences, University of Muhammadiyah Malang with research permit number E.5.d/134/PSIK/FIKES-UMM/II/2024. This research has also obtained approval or informed consent from all respondents and has received permission from the village head of the research area.

RESULTS

Respondent Characteristics

From the results of the research conducted, data were obtained in group 01 that 8 people (32%) participated in Gastroenteritis health education activities at the age 2330 years, 10 people (40%) at the age of 31-40years, 4 people aged 41-50 years (16%), and aged >50 years 3 people (12%). Regarding gender characteristics, all respondents were women (100%). In terms of educational characteristics, 3 people were elementary school educated (12%), 19 people were junior/high school educated (76%), and 3 people were college educated (12%). In terms of job characteristics, it was found that 1 person worked as a private employee (4%), 3 people worked as self-employee (12%), 6 people worked as traders (24%), and 15 people worked as housewives (60%). Based on the characteristics of many family members in one house, it was found that the number of family members was 2-4 people in one house, 21 people (84%) and the number of family members >5 people in one house was 4 people (16%).

From the results of the research conducted, data was obtained in group 02 that respondents who participated in Gastroenteritis health education activities were aged 21-30 years 1 person (4%), 31-40 years 4 people (16%), aged 41-50 years 14 people (56%), and age >50 years 6 people (24%). Regarding gender characteristics, all respondents were women (100%). In terms of educational characteristics, it was found that 8 people had an elementary school education level (32%), 11 people had a junior/high school education level (44%), and 6 people had a tertiary education level (24%). In terms of job characteristics, it was found that 1 person was unemployed (4%), 4 people were private employees (16%), 1 person was employed as a self-employed person (4%), 2 people were employed as traders (8%), 2 people were employed as laborers. people (8%), and 15 people (60%) work as housewives. Based on the characteristics of many family members in one house, it was found that the number of family members was 2-4 people in one house, 24 people (96%), and the number of family members >5 people in one house was 1 person (4%).

Frequency distribution of knowledge level after being given health education about Gastroenteritis using the lecture method and WhatsApp bot media

The results of the post-test data were obtained after being given health education about gastroenteritis 36 ng WhatsApp bot media, 9 people (36%) had a sufficient level of knowledge and 16 people (64%) had a good level of knowledge. While the post-test knowledge data for group 02 using lecture media, the results showed that 38 overall knowledge level was good for 25 people (100%).

Table 1. Frequency Distribution of Respondent Characteristics Based on Age of Group 01 and Group 02 (N=25)

Group 02 (11–22)							
Group 0	Group 02						
Characteristics	(f)	(%)	Characterist 24	(f)	(%)		
Age			Age				
<20 years	0	0	26<20 years	0	0		
21-30 years old	8	32	21-30 years old	1	4		
31-40 years old	10	40	31-40 years old	4	16		
41-50 years old	4	16	41-50 years old	14	56		
>50 years old	3	12	>50 years old	6	24		
Total	25	100	Total	25	100		
Educatio	n		Education	ı			
Elementary school	3	12	Elementary school	8	32		
Middle/ High school	19	76	Middle/ High school	11	44		
College	3	12	College	6	24		
Total	25	100	Total	25	100		
Work			Work	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			
No work	0	0	No work	1	4		
Private employee	1	4	Private employee	4	16		
Self-employed	3	12	Self-employed	1	4		
Trader	6	24	Trader	2	8		
Laborer	2	8	Laborer	2	8		
Housewife	15	60	Housewife	15	60		
Total	25	100	Total	25	100		
Family Member in Same Home			Family Member in S	ame H	ome		
2-4 persons	21	84	2-4 persons	24	96		
>5 persons	4	16	>5 persons	1	4		
Total	25	100	Total	25	100		

Table 2. Frequency Distribution of Knowledge Level After Being Given Health Education Using WhatsApp Bot Media in Group 01 and Using the Lecture Method in Group 02 (N=25)

Group 01			Group 02				Comparison of Group 01 and Group 02	
Characteristics	(f)	(%)	Mean	Characteristics	(f)	(%)	Mean	Sig
Post-test				Post-test				
Good	16	64		Good	25	100		
Enough	9	36	21.00	Enough	0	0	30.00	0.001
Not enough	0	0	21.00	Not enough	0	0	30.00	0.001
Total	25	100		Total	25	100		

DISCUSSION

Characteristics of Respondents Regarding Mother knowledge

Based on the research results, it was found that the frequency distribution based on age was at most 31-50 years. According to Rita et al., (2020) state that age has a relationship with a person's knowledge.

Age is one factor that influences a person's strength and mindset. As we get older, our strengths, thinking patterns 60 knowledge also develop and increase. The level of education also influences a person's knowledge

because the higher the education, the better the attitude, motivation, and knowledge. Education can change a person's perspective on things, especially through information obtained through education (Padila, Andri, & Andrianto, 2023). Mother education can increase mother understanding of the prevention and treatment of gastroenteritis and maternal behavior can also influence family health so that it can reduce the death rate due to gastroenteritis in children (Momoh et al., 2022; Thiam et al., 2019).

Education level is one of the factors that influences individual knowledge and the results of this research show that the majority of subjects have educated up to middle school/high school. These results indicate that there is a relationship between education [47]:1 and the incidence of gastroenteritis. This research is in line with research by Bayomy et al., (2024); Yuniarti & Vinnata, (2020) which states that education will influence a person's knowledge of preventive measures for gastroenteritis. In other words, the more someone knows the causes, triggers, signs, symptoms, and prevention of gastroenteritis, the greater the chance of avoiding potential gastroenteritis triggers.

In this study, it was found that the majority of mothers' jobs were housewives (59%). For working mothers, the knowledge and information they obtain or find out about children's health and things that can cause gastroenteritis in children is obtained more easily than mothers who don't work (Marianna & Yuli, 2022). This is supported to research by Zulfita et al., (2022) which states that there is a relationship between housewives' knowledge and incidence of gastroenteritis in children (pvalue <0.05). The education level and income of working mothers are significantly related to mother's knowledge (Kaçan, Pallos, & Özkaya, 2022).

Mother's level of knowledge after being given Gastroenteritis education

The level of knowledge is divided into three categories, namely good, enough, and not enough. The description of the subjects' knowledge after being given health education in this study was that group 01 obtained a sufficient level of knowledge for 9 subjects (18%) and a good level for 16 subjects (32%), while group 02 obtained a good level of knowledge for 25 subjects (50%).

Knowledge is information combined between understanding and action behavior (Alrafiaah et al., 2022). Behavior is greatly influenced by knowledge. The level of knowledge is one of the factors that influences human behavior (Widyakusuma Putra & Manalu, 2020). Increasing knowledge will affect a person's attitude toward health problems that occur (Aris Widayati et al., 2021).

The effectiveness of providing Gastroenteritis education based on WhatsApp bot and Lectures

The data results show that group 02 which received health education using lecture media was better than group 01 which received education using WhatsApp bot media. From the results 13 he Mann-Whitney test in the two groups, the p-value was 0.001 or <0.05, which means there is a significant difference between providing education based on WhatsApp bot media and lectures.

Education the learning process is provided to expand knowledge which is expected [12] lead to implementing a healthy lifestyle. Health education aims to increase public knowledge and awareness to maintain and improve thei dealth (Siti Aisah, Suhartini Ismail, 2021). Health education aims to transform people's knowledge and habits by actively promoting optimal health (Ariyanto & Fatmawati, 2021). Providing education can be more effective if used in the media because it is a tool used in the communication process to convey messages to respondents through certain media. In research by Melo et al., (2022) stated that video and booklet educational media are effective media often used to prevent gastroenteritis.

WhatsApp bot is an automation on Whatsapp by processes incoming messages according to the program flow embedded in it (Dewi, 2022). By developing an effective autoresponder chatbot you can improve services in providing information to users and helping users solve problems Hartling et al., (2023).

Providing education using the lecture method is an educational approach that can significantly increase knowledge toward certain objectives by presenting learning material through oral narrative (Mulyani & Subandi, 2020). In this study, it was found that the knowledge of mothers who were given education through lectures was mostly at a good level of knowledge so lectures were an effective medium that could be used in providing gastroenteritis education. This is in line with research by Jannah, Salfarina, & Riskawaty, (2024) that lectures have an effective influence on increasing knowledge in providing gastroenteritis education.

Providing adequate information after education about prevention and appropriate treatment can help in anticipating and treating gastroenteritis in children at home (Alghadeer et al., 2021). Gastroenteritis education based on WhatsApp bot and lectures influenced mothers' knowledge as demonstrated by good pretest results after being provided with information on gastroenteritis-related material.

CONCLUSION

Providing Gastroenteritis health education using WhatsApp bot media and lectures given to mothers in Singosari district affects increasing mothers' knowledge in preventing Gastroenteritis in children. There is a significant difference between providing education based on WhatsApp bot media and lectures. Gastroenteritis education using lecture media is more effective than WhatsApp bot media in increasing mothers' knowledge. The follow-up plan for this research has not yet been carried out but it is hoped that for the follow-up plan, respondents will be evaluated once a month with researchers asking again about the mother's knowledge about gastroenteritis via WhatsApp. It is hoped that future researchers will be able to carry out research using the innovations in this research over a long time.

ACKNOWLEDGMENT

Special thanks to the mothers in Singosari District, Malang Regency who participated i 23 his research. Thanks go to expressed to the Faculty of Health Sciences, University of Muhammadiyah Malang, which

has supported the research that has been carried out.

40 DECLARATION OF CONFLICTING INTEREST

There is no conflicting interest on this study.

FUNDING

The authors received financial support for manuscript research from the supervisor.

AUTHOR CONTRIBUTION

Sonia Puspita Ar Royo: Took responsibility for data analysis, interpretation, discussion of results, also writing, and editing manuscript.

Anis Ika Nur Rohmah: Contribution as a supervisor involved in planning and supervision in the completion of a manuscript.

Ika Rizki Anggraini: Contribution as a cosupervisor in reviewing the discussion of the final results of the manuscript.

Nur Melliza: Contribution as a co-supervisor in reviewing the discussion of the final results of the manuscript.

ORCID

Sonia Puspita Ar Royo:

https://orcid.org/0009-0007-0476-8972

Anis Ika Nur Rohmah:

https://orcid.org/0000-0001-8535-1851

Ika Rizki Anggraini:

https://orcid.org/0000-0001-7328-8939

Nur Melliza:

https://orcid.org/0000-0001-5533-2561

REFERENCES

Alrafiaah, AS, Albraikan, A., AlJaafari, A., AlAbbad, A., Alfehaid, H., Alqueflie, S., & Omair, A. (2022). Assessment of Maternal Knowledge and Practices

- Regarding Acute Diarrheal Illnesses in Children in Saudi Arabia: A Tertiary Care Center Survey. Cureus, 14(12). https://doi.org/10.775
- Ariani, D. Praise. (2022). Effectiveness of Counseling on the Level of Knowledge and Attitudes of Mothers of Toddlers Regarding Diarrhea Prevention. Proceedings of the National Seminar, 70–77.
- Arianti, M., & Jaya, H. (2022). Socialization and education on the importance of health knowledge about diarrhea in children and adults in the Tanjung Gading area, Bandar Lampung C 67

 Journal of Community Service, 1(1), 41–48.

 https://doi.org/10.59030/jpmbd.v1i1.1
- Aris Widayati, YW, Christasani3, PD, & Titien Siwi Hartayu2. (2021). Health Protocol Education Among Students. 4(3), 423– 428.
- Ariyant 19 A., & Fatmawati, TY (2021).

 Education on the Prevention of Diarrhea in Children in the Dasawisma Group, Recognize Lower Acid Subdistrict. Jour 35 of Community Health Greetings (JSSM), 2(2), 13–18. https://doi.org/10.22437/jssm.v2i2.136
- Arsi, R., Afdhal, F., & Fatrida, D. (2021).

 Education for a Healthy and
 Prosperous Community (EMaSS):
 Journal of Community Service. Journal
 of Community Service, 3(2), 1–4.
- Bayomy, HE, Almatrafi, HM, Alenazi, SF, Madallah S. Almatrafi, R., Alenezi, M., & Alanazi, WA (2024). Knowledge and Behavioral Practice of Mothers Regarding Childhood Diarrhea in A City, Saudi Arabia. Cureus, 16(2), 17.
- https://doi.org/10.7759/cureus.54221
 Dewi, RDC (2022). Early Warning System
 (Use of Wh 21 app Bot in the Health
 Sector). Bureaucracy Journal*;

- Indonesian Journal of Law and Social-Political Governance, 3(1), 10–19. https://doi.org/10.53363/bureau.v3i1.153
- Doris, A. (2021). Nursing Care for Children with a Diagnosis of Gastroenteritis.

 48 rebral Medika Scientific Journal, 1(1).
 - https://doi.org/10.53475/jicm.v1i1.62
- Febiyana, SC, Saputri, MA, Sari, PL, & Sari, WR (2024). Increasing Mothers' Knowledge about Making Tempe Porridge as an Effort to Treat Diarrhea in Toddlers in Ploso Village. 2(1).
- Giansanti, D. (2020). WhatsApp in mHealth: an overview on of the potentialities and opportunities in medical imaging.

 MHealth, 6(3), 1–5.

 https://doi.org/10.21037/mhealth.2019
- Hartling, L., Elliott, S. A., Munan, M., & Scott, S. D. (2023). Web-Based Knowledge Translation Tool About Pediatric Acute Gastroenteritis for Parents: Pilot Randomized Controlled Trial. JMIR Formative Research, 7. https://doi.org/10.2196/45276
- Herlina Alvianti Ningsih, Yulia Wardita, T. F. (2021). The Relationship between Providing Complementary Food for Breast Milk (MPASI) and the Occurrence of Diarrhea in Babies Before the Age of 6 Months in Pasean District. 4(1), 7–9.
- Heydarian, F., E, B., S.H., B., & M, H. (2019).

 Gastroenteritis Related Seizure with or without Fever: Comparison of Clinical Features and Serum Sodium Level How. 353900(5865), 47–52.
- Jannah, R., Salfarina, AL, & Riskawaty, HM (2024). Family education in preventing diarrhea in toddlers. Community Development Journal, 5(1), 355–359.
- Khoiriyah, Widjajanegara, H., & Sjafei, F. (2021). Scoping Review: Effectiveness of Giving Synbiotics to Pediatric Patients with Acute Gastroenteritis.

Proceedings of Medicine, 7(1), 783–794

Komara, IMAN, Jayadi, IPOK, Jayanti, NLPA,
Triyasa, P., M 34 gala, AK, & Sutisna,
P. (2020). The relationship between the
level of maternal knowledge about
preventing diarrhea and the incidence
of diarrhea in toddlers in Pemecutan
Kelod Village, Denpasa Bali.
Medical Science Digest, 11(3), 1247–
1251.

https://doi.org/10.15562/ism.v11i3.67

Lisa Hartling, P., Sarah A Elliott, P., Matthew
Muna 30 Ms., & Shannon D Scott, RN,
P. (2021). Assessment of Saudi
Mother's Knowledge and Attitudes
Towards Childhood Diarrhea and Its
Management. 25(2).

Manji, K., Hanefeld, J., Vearey, J., Walls, H., & De Gruchy, T. (2021). Using WhatsApp messenger for health systems research: A scoping review of available literature. Health Policy and Planning, 36(5), 774–789. https://doi.org/10.1093/heapol/czab024

Malang City Health Service. (2022). Malang City in 2021. *Malang City Health Service*, 45, 1–226.

Marianna, S., & Yuli, Utami. (2022). The level of maternal knowledge in serving formula milk influences the incidence of diarrhea in children aged 6-24 months. PROFESSION (Professional Islam), 20(1), 96–108. Retrieved from journals.itspku.ac.id

Melo, ESJ, de Oliveira, BSB, Melo, FM de S., da Silva, MJN, de Oliveira, 20 KL, Bezerra, JC, ... Barbosa, LP (2022). Audiovisual and printed technology to prevent childhood diarrhea: A clinical trial. Public Health Nursing, 39(2), 423–430.

https://doi.org/10.1111/phn.12962 Momoh, FE, Olufela, OE, Adejimi, AA, Roberts, AA, Oluwole, EO, Ayankogbe, OO, & Onajole, AT (2022). Mothers' knowledge, attitude and home management of diarrhea among children under five years old in Lagos, Nigeria. African Journal of Primary Health Care and Fan Medicine, 14(1), 1–10. https://doi.org/10.4102/phcfm.v14i1.3

Mulyani, S., & Subandi, A. (2020).

Effectiveness of Health Education through Periodic WhatsApp Reminder Groups Using the Lecture Method for Providing Exclusive Breastfeeding to Mothers After Cesarean Sections.

Jambi University 33 entific Journal of Applied Sciences JIITTUJI, 4(2), 187–203.

https://doi.org/10.22437/jiituj.v4i2.11 607

Mutmawardina, Harpiana Rahman, Farihah Muhsanah, Andi Asrina, & Nurul Likmah B. (2023). Differences in Lecture Methods and Audiovisual Media on Knowledge of Hand Washing with Soap at SD Inpres Binanga 3 Mamuju, West Sulwesi Province in 2022. Window of Public Health Journal, 4(2), 274–285. https://doi.org/10.33096/woph.v4i2.74

Nari, J. (2019). Nursing Care for Children with Acute Gastroenteritis to Fulfill Fluid and Electrolyte Needs in the Children's Room at RSUD Dr. M. Haulussy. 4(3), 131–136.

Padila, P., Andri, J., & Andrianto, MB (2023).

10 leo and Booklet Media Education on Parents' Motivation and Attitudes in

19 ing for Toddlers with Gastroenteritis. Journal of Telenursing (JOTING), 5(1), 720–731.

https://doi.org/10.31539/joting.v5i1.58

Purnamasari, T., & 14 tri, A. (2023).

Management of Health Education on

Mother's Knowledge in Efforts to

- Prevent Diarrhea in Children Under Five (0-5 Years) in 2022. Journal of Health, 9(1), 65–73.
- Qori Nurul Isnaini, EF (2021). Nursing in Gastroenteritis Patients With Disorders of, 15.
- Rachman, DD, & Nurjanah, D. (2019). Ontology Model Recommendation System for Providing Diet Recommendations for Diabetes Sufferers. E-Proceedings of 9569-9584. grgineering, 6(2), Retrieved from https://openlibrarypublications.telkom university.ac.id/index.php/engineering /article/viewFile/10002/9858
- Rahmawati, NE, & Arifiyanti, AA (2020).

 Diagnosis of Inflammatory Arthritis

 14 ng the Certainty Factor Method.

 SATIN Information Science And
 Technology, 3(1), 42–47.

 https://doi.org/10.33372/str
- Rimbawati, Y., & Surahman, A. (2019). The
 Relationship between Environmental
 Sanitation and the Incidence of
 Diarrhea in Toddlers. Journal of
 'Aisyiyah Medika, 4, 189–198.
 https://doi.org/10.36729/jam.v4i0.337
- Rita, R., Amir, N., & Suhardi, D. (2020). The
 Relationship Between Knowledge of
 Parents of Patients and Handling
 Diarrhea in the Children's R 10 n at
 Jayapura Regional Hospital. Sentani
 Nursing Journal, 3(2), 44–53.
 https://doi.org/10.52646/snj.v3i2.44
- Salam, MA us, Oyekwe, 15. Ghani, SA, & Choudhury, RI (2021). How can WhatsApp® facilitate the future of medical education and clinical practice? BMC Medical Education, 21(1), 1–4. https://doi.org/10.1186/s12909-020-02440-7
- Siahaan, YF, Aritonang, EY, & Ashar, T. (2020). Relationship between Infant Feeding Practices and the Incidence of Gastroenteritis. Journal of the

- Indonesian 49 tional Nurses
 Association (JPPNI), 4(1), 1.
 https://doi.org/10.32419/jppni.v4i1.17
- Siti Aisah, Suhartini Ismail, AM (2021). Health
 Education Using Animation Video
 Media: Scoping Revie 10 Indonesian
 Nursing Journal, 5(1), 641–655.
 https://doi.org/10.32584/jpi.v5i1.926
- Susilowati, IT, & Suwarni, S. (2023).

 Prevention and Management of
 Diarrhea and Vomiting through
 Education on Clean Living Behavior
 and the Use of Herbal Ingredients.

 PARAHITA: Jour 68 of Community
 Service, 3(2), 55–64.

 https://doi.org/10.25008/parahita.v3i2.
- Thiam, S., Sy, I., Schindler, C., Niang-Diène, A., Faye, O., Utzinger, J., & Cissé, G. (2019). Knowledge and practices of mothers and caregivers on diarrheal management among under 5-year-old children in a medium-sized town of Senegal. Acta Tropica, 194(February), 155–164. https://doi.org/10.1016/j.actatropica.2 019.03.013
- Umma, N., Sari, NPD, Afifah, RN, Baharudin, MY, Pribadi, P., Santoso, SB, & Latifah, E. (2024). Drug Consultagen Prototype Based WhatsApp Bot (Vol. 2022). Atlantis Press SARL. https://doi.org/10.2991/978-2-38476-118-0_137
- Widyakusuma Putra, YI, & Manalu, NV (2020). Level of Knowledge and Citizen Behavior in Implementing Health Protocols in the New 22 rmal Corona Pandemic Period. Coping: Community of Publishing in Nursing, 8(4), 366. https://doi.org/10.24843/coping.2020. v08.i04.p04
- Yona, CC, & Iriani, R. (2022). Persada Husada Indonesia Journal Nursing Care of Clients Who Experience Lack of Body

Fluid Volume with Gastroenteritis at Budhi And Regional General Hospital, Jakarta Nursing Care of Clients Who Experience Lack of Body Fluid Volume with Gastroenter. 9(32), 46–54.

Yüksel Kaçan, C., Palloş, A., & Özkaya, G. (2022). Examining knowledge and traditional practices of mothers with children under five in Turkey on diarrhea according to education levels.

Annals of Medicine, 54(1), 674–682. https://doi.org/10.1080/07853890.202 2.2044508

Yuniarti E., & Vinnata, NN (2020).

Relationship of Education and Knowledge of Formula Milk Feeding to Dia Glea in Children Aged 0 – 2

Years Relationship of Education and Knowledge of Formula Milk He eding for. CITRA DELIMA: Scientific Journal of STIKES Citra Delima

Bangka Belitung, 4(1), 7–11.

Zulfita, A., Sari, NP, Wardani 17, Yulianto, B., & Hayana, H. (2022). The Relationship Between Housewives' Personal Hygiene and Basic Sanitation Facilities and the Incidence of Diarrhea in Toddlers in the Sapta Taruna Center Work Area in 2021. Public

Health Media, 2(1), 151–161. https://doi.org/10.25311/kesmas.vol2.i ss1.512

Cite this article as: Ar Royo, S. P., et al. (2024). Effectiveness of Whatsapp Bot-based E 33 ational Media and Lectures on Increasing Mother's Knowledge about Gastroenteritis in Children. Nurse and Health: Jurnal Keperawatan, 13 (2), 19-29. https://doi.org/10.36720/nhjk.v13i2.663

EFFECTIVENESS OF WHATSAPP BOT-BASED EDUCATIONAL MEDIA AND LECTURES ON INCREASING MOTHER'S KNOWLEDGE ABOUT GASTROENTERITIS IN CHILDREN

ORIG	INALITY REPORT	
	3% ARITY INDEX	
PRIM	ARY SOURCES	
1	media.neliti.com Internet	70 words — 1 %
2	jurnal.fkm.umi.ac.id Internet	65 words — 1 %
3	repository.unhas.ac.id Internet	50 words — 1 %
4	mpu.med-expert.com.ua Internet	45 words — 1 %
5	eprints.whiterose.ac.uk	39 words — 1 %
6	repo.undiksha.ac.id Internet	39 words — 1 %
7	jptcp.com Internet	37 words — 1 %
8	journal-mandiracendikia.com Internet	36 words — 1 %
9	jurnalilmiah.stikescitradelima.ac.id	36 words — 1 %

10	journal.ipm2kpe.or.id Internet	35 words — 1 %
11	jurnal.stikesphi.ac.id Internet	35 words — 1 %
12	Carsten Posovszky, Stephan Buderus, Johannes Huebner, Rüdiger Adam et al. "S2k Leitlinie akute infektiöse Gastroenteritis im Säuglings-, Kindes- ur Jugendalter – Update 2024", Zeitschrift für Gastroe 2024 Crossref	
13	Eva Dwi Ramayanti, Indah Jayani, Satria Eureka Nurseskasatmata, Arif Nurma Etika, Wiwin Sulistyawati. "PENGARUH PENDIDIKAN KESEHATAN MENGGUNAKAN MEDIA VIDEO INFORMATIF TERH PERUBAHAN TINGKAT STRES PADA ODHA DI MASA COVID-19", Nursing Sciences Journal, 2023 Crossref	ADAP
14	Andi Elis, Rohani Mustari, Marlina Marlina. "The Ro of Presenting Exclusive Breastfeeding for The Prevention of Stunting Based on The Culture of Tu Sipulung", STRADA Jurnal Ilmiah Kesehatan, 2020 Crossref	
15	4d.tecnocampus.cat Internet	27 words — < 1%
16	thesis.unipd.it Internet	27 words — < 1%
17	journal.uinjkt.ac.id	26 words — < 1%

18	ojs.uho.ac.id Internet	26 words — < 1%
19	repository.universitas-bth.ac.id	24 words — < 1 %
20	Emanuella Silva Joventino Melo, Brena Shellem Bessa de Oliveira, Francisca Mayra de Sousa Melo, Maria Jocelane Nascimento da Silva et al. "A and printed technology to prevent childhood diar trial", Public Health Nursing, 2021 Crossref	
21	bureaucracy.gapenas-publisher.org	20 words — < 1 %
22	repository.binausadabali.ac.id Internet	20 words — < 1 %
23	eprints.umm.ac.id Internet	17 words — < 1 %
24	repository.unp.ac.id Internet	17 words — < 1 %
25	elearning.medistra.ac.id Internet	16 words — < 1%
26	en.hpchsu.ru Internet	16 words — < 1%
27	eprints.mums.ac.ir Internet	15 words — < 1%
28	journal.formosapublisher.org	15 words — < 1%

29	jurnal.uisu.ac.id Internet	15 words — <	1%
30	www.eimj.org Internet	15 words — <	1%
31	wjgnet.com Internet	14 words — <	1%
32	ejournal.umm.ac.id Internet	13 words — <	1%
33	Ema Pristi Yunita, Rayfika Nokha Kusuma Wardani, Bambang Sidharta. "Correlation between knowledge level, side effect severity, far and antiretroviral therapy adherence in HIV/AIDS Greater Malang, East Java, Indonesia", Pharmacia Crossref	patients in	1%
34	Ritzkal Ritzkal, Kodarsyah, Puspa Putri Amalia, Wildan Mahmud, Ade Hendri Hendrawan, Bayu Adhi Prakoso, Indra Riawan. "Security Vulnerabilit and Recommendations for Open Media Vault Clo Raspberry Pi", Ingénierie des systèmes d informa Crossref	ud Server on	1%
35	jurnal.padangtekno.com Internet	12 words — <	1%
36	jurnal.stikeshusadajombang.ac.id Internet	12 words — <	1%
37			0.1
	jurnal.umt.ac.id Internet	12 words — <	1%

39	www.jurnal.poltekkesgorontalo.ac.id	12 words — <	1%
40	Intansari Nurjannah. ""SU JOK" THERAPY AND SCLEROLOGY PROFILE MONITORING FOR MANAGING CHEST PAIN AT HOME WHILE AVOID ADMISSION DURING THE COVID-19 PANDEMIC: STUDY", Belitung Nursing Journal, 2020 Crossref		1%
41	formative.jmir.org Internet	11 words — <	1%
42	pjbt.org Internet	11 words — <	1%
43	www.abdn.ac.uk Internet	11 words — <	1%
44	www.e-journal.unair.ac.id Internet	11 words — <	1%
45	Dina A. Alkhozaim, Sanaa N. Al-Haj Ali, Ra'fat I. Farah. "Levels and correlates of knowledge of teething among Saudi Arabian families", PeerJ, 20 Crossref	10 words — <	1%
46	ejournal.ipinternasional.com Internet	10 words — <	1%
47	j-innovative.org Internet	10 words — <	1%
48	jurnal.globalhealthsciencegroup.com Internet	10 words — <	1%

49	repository.penerbiteureka.com Internet	10 words —	<	1	%
50	www.isainsmedis.id Internet	10 words —	<	1	%
51	Akma Listiana, Novi Eniastina Jasa. "The Influence Of Health Education Using Video Education Media On Increasing Adolescent Women's Knowledge Ak JKM (Jurnal Kebidanan Malahayati), 2023 Crossref			1	%
52	humaniora.journal.ugm.ac.id Internet	9 words —	<	1	%
53	icudspace.icuzambia.net Internet	9 words —	<	1	%
54	journal.stikesdrsoebandi.ac.id Internet	9 words —	<	1	%
55	journalajmah.com Internet	9 words —	<	1	%
56	jurnal.unitri.ac.id Internet	9 words —	<	1	%
57	pjmd.zu.edu.pk Internet	9 words —	<	1	%
58	www.frontiersin.org Internet	9 words —	<	1	%
59	Elliot Mbunge, Maureen Nokuthula Sibiya, Sam Takavarasha, Richard C Millham et al. "Implementation of ensemble machine learning c	8 words — lassifiers to	<	1	%

predict diarrhoea with SMOTEENN, SMOTE, and SMOTETomek class imbalance approaches", 2023 Conference on Information Communications Technology and Society (ICTAS), 2023

Crossref

Sabtria Winda Sari, Erika ., Awaluddin ., Winda Ulan 8 words — <1% Sari. "The Effect of Warm Ginger Water Foot Bath on Blood Pressure Reduction in Hypertensive Elderly at Sialang Baru", KnE Medicine, 2023

Crossref

Sri Haryani, Yusniarita Yusniarita, Yanti Sutriyanti, $_{8 \text{ words}} - < 1\%$ Derison Marsinova Bakara. "Knowledge and Attitude of Cadres in the Prevention and Treatment of Diarrhea At Home Among Mothers of Toddlers", JOSING: Journal of Nursing and Health, 2024

Crossref

- assets-eu.researchsquare.com 8 words < 1%
- ijor.co.uk
 _{Internet}
 8 words < 1 %
- journals.stikim.ac.id
 _{Internet}
 8 words < 1%
- repository.canterbury.ac.uk 8 words < 1%
- www.e-journal.poltekkesjogja.ac.id
 8 words < 1 %
- www.jiemar.org
 Internet

 8 words < 1 %
- 68 www.sciencepubco.com

9 words	<	1	%
8 words -	_ `		, 0

- www.trustedhealthproducts.com
- 8 words < 1%

 7 words < 1% 70 Novi Eka Fitrah, Meri Neherta, Ira Mulya Sari. "Gambaran Pengetahuan dan Sikap Ibu dalam Pemberian ASI Eksklusif sebagai Upaya Pencegahan Diare pada Anak Balita", Journal of Telenursing (JOTING), 2023 Crossref

OFF

EXCLUDE BIBLIOGRAPHY OFF OFF